

FULL TEXT LINKS



Observational Study J Pediatr Adolesc Gynecol. 2020 Oct;33(5):550-554.

doi: 10.1016/j.jpag.2020.05.010. Epub 2020 Jun 11.

# The Association between Immediate Postpartum Etonogestrel Implants and Positive Postpartum Depression Screens in Adolescents and Young Adults

Erin Drake <sup>1</sup>, Kira Grush <sup>2</sup>, Jeanelle Sheeder <sup>2</sup>, Kristina Tocce <sup>3</sup>

Affiliations

PMID: 32535218 DOI: 10.1016/j.jpag.2020.05.010

## Abstract

**Objective:** We aimed to compare rates of positive postpartum depression screens at 6 weeks postpartum among adolescents and young adults (AYA) initiating immediate postpartum contraceptive implants and those initiating other methods.

**Design:** Through a retrospective observational design, we collected data on demographics, reproductive history, prenatal and postnatal depression, and postpartum contraception.

**Setting:** Patients participating in an AYA prenatal-postnatal program were eligible for inclusion.

**Participants:** A total of 497 patients were enrolled between January 2013 and December 2016. The median age was 19 years (range 13-22 years); 86% were primiparous, 50% were Latina, 24% were black, and 16% were white; 34% initiated immediate postpartum implants (n = 169).

**Intervention:** Those initiating a contraceptive implant within the first 14 days postpartum were included in the intervention group.

**Main outcome measure:** We compared rates of positive Edinburgh Postpartum Depression Scales (EDPS) (scores  $\geq 10$ ) in AYA initiating immediate postpartum implants and those initiating other contraceptive methods.

**Results:** The AYA initiating immediate postpartum implants were similar to the rest of the cohort in baseline characteristics, aside from an increased rate of preterm births among the intervention group (19.4% vs 12.1%;  $P = .03$ ). Prenatally, 14% had an elevated Center for Epidemiologic Studies Depression Scale (CES-D) scores (11.5% immediate postpartum implants vs 15.4% comparison,  $P = .25$ ). At 6 weeks postpartum, 7.6% had a positive postpartum depression screen; this rate was significantly lower for those initiating immediate postpartum implants compared to those choosing other methods (4.1% vs 9.5%,  $P = .04$ ).

**Conclusions:** Providers should continue to encourage AYA to choose whichever highly effective contraceptive method they prefer for postpartum use.

**Keywords:** Adolescent; Immediate postpartum implant; Long-acting reversible contraception; Postpartum depression.

Copyright © 2020 North American Society for Pediatric and Adolescent Gynecology. Published by Elsevier Inc. All rights reserved.

## Related information

MedGen