

Platelet/Lymphocyte Ratio: A Potential Biomarker for Disease Activity in Ankylosing Spondylitis

EJ Kim, BA¹; R Sen, MD MS^{1,2}; ES Manning, MD^{1,2}; ER Anderson, MD¹; KD Maier, MD¹; E Cheng, BA²; L Caplan, MD PHD^{1,2}
¹ University of Colorado School of Medicine, Aurora, CO, ² Rocky Mountain Regional Veterans Affairs Medical Center, Aurora, CO



Study Aim

Evaluate the utility of the neutrophil/lymphocyte ratio and platelet/lymphocyte ratio as biomarkers for Ankylosing Spondylitis disease activity and severity by examining their association with c-Reactive Protein and sacroiliac joint damage scores.

Background

Ankylosing spondylitis: a chronic inflammatory disease involving the sacroiliac joint (SIJ) and the spine → over time, joints can be involved with erosions or fusion that can be debilitating and affect the quality of life.

C-Reactive Protein (CRP): low specificity and sensitivity, but is the most commonly used biomarker for disease activity.

Neutrophil/lymphocyte ratio (NLR) and Platelet/lymphocyte ratio (PLR): shown in prior studies to be associated with disease activity and severity for inflammatory diseases such as malignancies and ulcerative colitis. A high NLR and PLR reflect inflammation.

Results

Demographics			Multiple linear regression (n=42)						
	N	Mean	Dependent variable	Independent variable	Coefficient	P value	95% CI lower	95% CI upper	R ²
Age (years)	42	51.9	SIJ mNY total score						
Married, %	41	56.1		PLR	0.010	0.064	-0.001	0.0215	0.1015
White, %	42	78.6		NLR	-0.137	0.506	-0.548	0.275	
Black, %	42	9.52							
HLA_B27 (+), %	42	76.2	Simple linear regression (n=37)						
Male, %	42	97.6	Dependent variable	Independent variable	Coefficient	P value	95% CI lower	95% CI upper	R ²
Current smoker, %	42	24.3	CRP	PLR	0.244	<0.001	0.122	0.366	0.3205
Former smoker, %	41	39.0		NLR	2.304	0.403	-3.219	7.827	0.0201
Never smoker, %	41	36.6							

Methods

Population characteristics	• Program to Understand Long Term Outcomes of Spondyloarthritis (PULSAR) registry at Rocky Mountain Regional VA Hospital
SIJ radiograph scored for sacroiliitis	• scored using the modified New York criteria (mNY). • excluded those who did not meet mNY criteria
Obtain CBC and CRP at time of Xray (+/- 6 mo)	
Data analysis	• Determined relationship between NLR, PLR, CRP and mNY score using regression techniques in STATA (v13) statistical package

Conclusion

- No statistically significant correlation was found between mNY radiograph scores and the NLR nor the PLR.
- CRP concentration was associated with the PLR.

Future Directions

- Validate these results in other populations
- Explore the relationship of PLR with patient reported outcomes
- Determine if PLR is associated with findings from other imaging modalities.

Acknowledgements

This research project was funded by the Schweppe Donation. No disclosures or conflicts of interest.