

BACTERIAL MENINGITIS AND PEDIATRIC SENSORINEURAL HEARING LOSS IN THE POST-PCV13 ERA.



University of Colorado
Anschutz Medical Campus

Shilpa Tummala, BA¹, Kaitlyn Tholen, BS², Maxene Meier^{1,3}, Brian Herrmann, MD^{1,2}, Juri Boguniewicz, MD^{1,4}, Sarah Gitomer, MD^{1,2}

¹University of Colorado Anschutz School of Medicine, 13001 East 17th Place, Aurora, CO 80045

²Children’s Hospital of Colorado (CHCO), Anschutz Medical Campus, Department of Otolaryngology—Pediatric, 13123 East 16th Avenue, Aurora, CO 80045

³Center for Research Outcomes in Children’s Surgery, Center for Children’s Surgery, University of Colorado Anschutz Medical Campus, Aurora, CO 80045

⁴Children’s Hospital of Colorado (CHCO), Anschutz Medical Campus (CHCO), Department of Infectious Disease—Pediatrics, 13123 East 16th Avenue, Aurora, CO 80045

The authors do not have any relevant financial conflicts to disclose.

ABSTRACT

INTRODUCTION: Bacterial meningitis is the most common cause of postnatally-acquired sensorineural hearing loss (SNHL). Prognostic associations are lacking for pediatric SNHL due to *Streptococcus pneumoniae* bacterial meningitis after the introduction of updated pneumococcal vaccine (PCV13) in 2010. **OBJECTIVES: 1)** To assess clinical presentation of meningitis based on bacterial etiology in children and identify associations with post-meningitic SNHL in children after 2010. **2)** To describe rates of *S. pneumoniae*-associated post-meningitic SNHL based on three time periods: pre-PCV, post-PCV7 and post-PCV13. **METHODS:** A retrospective review was performed for patients 18 years and younger diagnosed with meningitis after January 1, 2010. Patients were identified by history of positive CSF bacterial culture or FilmArray Meningitis/Encephalitis Panel (MEP) assay. Clinical data were stratified by bacterial etiology, analyzed for clinically relevant characteristics, and compared with previously reported rates of post-meningitic SNHL. **RESULTS:** In a cohort of 147 patients with positive CSF cultures, 91 (61.9%) met inclusion criteria. Seventy-one (mean age 21 months, 56% male) had audiograms after diagnosis and were divided into four subgroups based on bacterial etiology: *S. pneumoniae* (n=20), Group B streptococcus (GBS, n=25), *Haemophilus influenzae* (n=10), and Other (n=16). Of those with hearing evaluations, SNHL was reported in 15 patients (21%), most frequently in the *S. pneumoniae* population (n=8/19, 42%). Two (20%) *H. influenzae* patients and two (8%) GBS patient were found to have SNHL. Children in the PCV13 vaccination era had a similar rate of post-meningitic SNHL (42%) as historical pediatric cohorts in the pre-PCV vaccination time period (23.8%), and the PCV7 vaccination time period (35%) (Fisher’s exact, p=0.19). **CONCLUSION:** Despite advances in vaccine development for *S. pneumoniae*, SNHL remains a common long-term complication of this disease. Further research into predicting and preventing this outcome is necessary.

BACKGROUND

- Bacterial meningitis (BM) accounts for 60-90% of all pediatric sensorineural hearing loss (SNHL) cases and is associated with significant morbidity and mortality in children.
- The two most common causes of postnatally acquired SNHL cases are *H. influenza* type B (HiB) and *S. pneumoniae*.
- Vaccines against HiB and *S. pneumoniae* have dramatically reduced incidence of bacterial meningitis.
- However, the incidence of pediatric SNHL due to pneumococcal meningitis has remained stable at 28-35% between the years of 2007-2013.
- Hearing loss has significant effects on childhood development, contributing to delays in language development and academic performance.
- No prior studies have compared bacterial meningitis and hearing outcomes from children in the pre-PCV, post-PCV7 and post-PCV13 eras.

OBJECTIVES

- To present a pediatric cohort of patients with bacterial meningitis and SNHL in the post-PCV13 era
- To compare historical data about bacterial meningitis and SNHL from the pre-PCV and post-PCV7 eras with a post-PCV13 pediatric cohort

METHODS AND MATERIALS


- Institutional review board approval was obtained.
- Chart review of patients with spontaneous bacterial meningitis from 2010-2020
- **Inclusion criteria:** CHCO patients diagnosed with spontaneous bacterial meningitis on or after January 1, 2010. Age 18 years or younger at time of diagnosis. Diagnosis must be made by positive CSF cultures.
- **Exclusion criteria:** Patients with known underlying immunodeficiencies, on biologic agents such as monoclonal antibodies or with bacterial meningitis related to trauma, a shunt infection or a neurological procedure. Patients without culture or PCR-confirmed meningitis due to bacterial etiology.
- Data was compared with previously reported rates of post-meningitic SNHL.

RESULTS

Demographics

- 147 patients between 2010-2020
 - 91 (61.9%) met inclusion criteria
 - Mean age at diagnosis: 23.13 mos
- A total of 71 patients had post-meningitic audiograms available


Meningitis by Bacterial Etiology



Outcomes

- 19 total patients exhibited post-meningitic hearing loss (HL)
- 2 patients had HL prior to onset of meningitis 5 patients had conductive hearing loss
- 2 patients had mixed HL, both of whom also had SNHL in the other ear
- Pneumococcal meningitis was the most common cause of post-meningitic SNHL with six affected patients (42%)

SNHL in Post-Pneumococcal Meningitis Patients



	H. flu (N=10)	Other (N=16)	GBS (N=25)	S. pneumoniae (N=20)	Total (N=71)
Age (months)	64.36	18.11	1.15	25.47	20.73
Gender					
Male	6 (60%)	12 (75%)	8 (32%)	14 (70%)	40 (56%)
Female	4 (40%)	4 (25%)	17 (68%)	6 (30%)	31 (44%)
Ethnicity					
Hispanic or Latino	1 (13%)	5 (33%)	5 (21%)	6 (33%)	17 (25%)
Not Hispanic or Latino	7 (88%)	10 (67%)	19 (79%)	13 (67%)	49 (75%)
Race					
White	5 (50%)	11 (69%)	21 (84%)	15 (75%)	52 (74%)
Black/African American	3 (30%)	3 (19%)	0 (0%)	0 (0%)	6 (9%)
Asian	0 (0%)	0 (0%)	1 (4%)	0 (0%)	1 (1%)
More than one race	0 (0%)	1 (6%)	1 (4%)	1 (5%)	3 (3%)
Other	0 (0%)	1 (6%)	2 (8%)	3 (15%)	6 (9%)
Unknown/Not reported	2 (20%)	0 (0%)	0 (0%)	1 (5%)	3 (3%)

Table 1. Demographics and Clinical Characteristics stratified by Bacterial Etiology

Table 2. Outcomes for patients with bacterial meningitis stratified by bacterial etiology

	H. flu (N=10)	Other (N=16)	GBS (N=25)	S. pneumoniae (N=20)	Total (N=71)
Hearing Loss?					
Yes	4 (40%)	4* (27%)	2 (8%)	9* (47%)	19 (28%)
No	6 (60%)	11 (73%)	23 (92%)	10 (53%)	50 (72%)
Deceased?					
Yes	0 (0%)	1 (6%)	0 (0%)	0 (0%)	1 (1%)
No	10 (100%)	15 (94%)	25 (100%)	20 (100%)	70 (99%)

*1 patient was found to have hearing loss prior to bacterial meningitis diagnosis

DISCUSSION

- We present a novel analysis of post-meningitis SNHL in a pediatric cohort after the introduction of the PCV13 vaccine in 2010.
- Prior studies from the pre-PCV7 and PCV7-vaccinated eras have reported similar rates of post-meningitic SNHL as our pediatric cohort.
- Our results indicate a need for further exploration of predictive markers in the pathogenesis of post-meningitic SNHL, particularly in pediatric pneumococcal populations.

CONCLUSION

- Advances in vaccine development have helped reduce incidence of bacterial meningitis in the US due to *H. influenzae* and *S. pneumoniae*.
- Rates of SNHL within post-pneumococcal meningitic populations remain constant in pediatric cohorts since the introduction of PCV7 and PCV13.
- Future directions include univariate testing and potential multivariate analysis if more patients are added that meet inclusion criteria.

Contact

Shilpa Tummala
Phone: 602-309-9702
Email: shilpa.tummala@cuanschutz.edu

References

1. Dodge PR, Davis H, Feigin RD, et al. Prospective evaluation of hearing impairment as a sequela of acute bacterial meningitis. *N Engl J Med.* 1984;311(14): 869-874.
2. Davis JM, Effenbein J, Schum R, Bentler RA. Effects of mild and moderate hearing impairments on language, educational and psychosocial behavior of children. *J Speech Hear Disord.* 1986;51(1): 53-62.
3. From the Centers for Disease Control and Prevention. Progress toward elimination of Haemophilus influenza type b invasive disease among infants and children – United States, 1998-2000. *JAMA.* 2002;287(17):2206-2207
4. Olarte L, Barson WJ, Barson RM, et al. Impact of the 13-Valent Pneumococcal Conjugate Vaccine on Pneumococcal Meningitis in US Children. *Clin Infect Dis.* 2015;61(5):767-775.
5. Richardson MP, Red A, Tarlow MJ, Rudd PT. Hearing loss during bacterial meningitis. *Arch Dis Child.* 1997;76(2): 134-138.
6. Wellman MB, Sommer DD, McKenna J. Sensorineural hearing loss in postmeningitic children. *Otol Neurotol.* 2003; 24(6): 907-912.
7. Woolley AL, Kirk KA, Neumann AM Jr, McWilliams SM, Murray J, Freind D, Wiatrak BJ. Risk factors for hearing loss from meningitis in children: the Children’s Hospital experience. *Arch Otolaryngol Head Neck Surg.* 1999 May;125(5):509-14. doi: 10.1001/archotol.125.5.509. PMID: 10326807.