

REVIEW

The power of zebrafish models for understanding the co-occurrence of craniofacial and limb disorders

Brittany T. Truong^{1,2} | Kristin B. Artinger² 

¹Human Medical Genetics & Genomics
Graduate Program, University of Colorado
Denver Anschutz Medical Campus, Aurora,
Colorado

²Department of Craniofacial Biology,
University of Colorado Denver Anschutz
Medical Campus, Aurora, Colorado

Correspondence

Kristin B. Artinger, Department of Craniofacial
Biology, University of Colorado Anschutz
Medical Campus, 12801 E. 17th Ave.,
MS8120, Aurora, CO 80045.
Email: kristin.arteringer@cuanschutz.edu

Funding information

Eunice Kennedy Shriver National Institute of
Child Health and Human Development
(NICHD), Grant/Award Numbers: NIH
1F31HD103368-01 (B. T. T.), NIH
1R03HD096320-01A1 (K. B. A.)

Summary

Craniofacial and limb defects are two of the most common congenital anomalies in the general population. Interestingly, these defects are not mutually exclusive. Many patients with craniofacial phenotypes, such as orofacial clefting and craniosynostosis, also present with limb defects, including polydactyly, syndactyly, brachydactyly, or ectrodactyly. The gene regulatory networks governing craniofacial and limb development initially seem distinct from one another, and yet these birth defects frequently occur together. Both developmental processes are highly conserved among vertebrates, and zebrafish have emerged as an advantageous model due to their high fecundity, relative ease of genetic manipulation, and transparency during development. Here we summarize studies that have used zebrafish models to study human syndromes that present with both craniofacial and limb phenotypes. We discuss the highly conserved processes of craniofacial and limb/fin development and describe recent zebrafish studies that have explored the function of genes associated with human syndromes with phenotypes in both structures. We attempt to identify commonalities between the two to help explain why craniofacial and limb anomalies often occur together.

KEYWORDS

craniofacial, human clinical genetics, limb, zebrafish model

1 | INTRODUCTION

Structural birth defects, including craniofacial and limb anomalies, affect 3% of newborns in the United States (Update on Overall Prevalence of Major Birth Defects, 2008). Human craniofacial defects occur in 1 in every 500–1,000 live births, with orofacial clefts being the most common (1:700) (Byvaltsev, Belykh, & Belykh, 2012; Global registry and database on craniofacial anomalies, 2001), while congenital limb disorders affect 1 in 1,000–2,000 newborns (Vasluian et al., 2013; Wilcox, Coulter, & Schmitz, 2015). Interestingly, these anomalies are not mutually exclusive. Many patients with craniofacial anomalies also present with upper and/or lower limb defects, such as syndactyly (digit fusion), ectrodactyly (missing digits), polydactyly (extra digits), and/or brachydactyly (shortening of the hands/feet). Syndromes that present with defects in both structures include Apert

syndrome (MIM #101200), Pfeiffer syndrome (MIM #101600), Roberts syndrome (MIM #268300), and Saethre-Chotzen syndrome (MIM #101400) among others (Table 1). This may be due to the synchronous timing of their development (Panthaki & Armstrong, 2003). Others have proposed that paired limbs/fins are evolved from gill or pharyngeal arch skeletal elements, components of which give rise to the craniofacial skeleton (Gegenbaur, 1878). These studies suggest that there is a deep homology between the two structures. Many published reviews have thoroughly discussed vertebrate craniofacial or limb development separately and described how genetic or environmental factors can lead to congenital defects (Mercader, 2007; Mork & Crump, 2015; Petit, Sears, & Ahituv, 2017; Szabo-Rogers, Smithers, Yakob, & Liu, 2010; Twigg & Wilkie, 2015; A. Zuniga, 2015); however, to our knowledge, no one has adequately explored how or why craniofacial and limb anomalies often occur together.

TABLE 1 Human congenital disorders with both craniofacial and limb defects modeled in zebrafish

Syndrome	Human gene	Inheritance pattern	Major human craniofacial/limb phenotypes	Zebrafish ortholog	Model	Zebrafish craniofacial phenotypes	Zebrafish pectoral fin phenotypes	Experimental assays	Reference
Achondroplasia (MIM #100800)	FGFR3 ^b	AD	Macrocephaly; prominent forehead; brachydactyly; trident hand; bowed legs	<i>fgfr3</i>	CRISPR-Cas9 LOF mutant	No phenotype	No phenotype	RT-qPCR; RNA ISH; skeletal staining	(Leerberg, Hopton, & Draper, 2019)
Acrofacial dysostosis, Cincinnati type (MIM #616462)	POLR1A	AD	Variable mandibulofacial dysostosis; micrognathia; upper/lower eyelid clefts; microcephaly; brachydactyly	<i>polr1a</i>	LOF mutant	Hypoplastic/deformed: Meckel's cartilage; ceratohyal; trabeculae Absent: Ethmoid plate; hyosymplectic cartilage; ceratobranchial arches; palatoquadrate	Hypoplastic: Pectoral fin ^d	RNA ISH; immunostaining; TUNEL assay; live-cell imaging; skeletal staining	(Weaver et al., 2015)
Acromelic frontonasal dysostosis (MIM #603671)	ZSWIM6	AD	Frontonasal dysplasia; nasal clefting; carp-shaped mouth; preaxial polydactyly; tibial hypoplasia/aplasia	<i>zswim6</i>	N/A	N/A	N/A	RNA-FISH	(Smith et al., 2014)
Alagille syndrome 1 (MIM #118450)	JAG1	AD	Broad, prominent forehead; deep, wide, low-set eyes; mandibular prognathia; brachydactyly	<i>jag1a/jag1b</i>	MO	Subtle midbrain and forebrain defects Hypoplastic: Ears Hypoplastic/deformed: Palatoquadrate; hyosymplectic cartilage; opercle bone Deformed: Meckel's cartilage; ceratohyal; hyosymplectic cartilage Ectopic cartilage near palatoquadrate and midline	N/A	RNA ISH; skeletal staining	(Lorent et al., 2004)
Alagille syndrome 2 (MIM #610205)	NOTCH2	AD	Broad, prominent forehead; deep, wide, low-set eyes; mandibular prognathia; brachydactyly	<i>notch2</i>	MO	N/A	N/A	RNA ISH	(Lorent et al., 2004)
Andersen-Tawil syndrome (MIM #170390)	KCNJ2	AD	Broad forehead; micrognathia; wide-set eyes; low-set ears; syndactyly; clinodactyly	<i>kcnj2a/kcnj2b</i>	DN mRNA OE	Deformed: Palatoquadrate; hyosymplectic cartilage; ceratohyal Wide-set eyes; protruding jaw	Hypoplastic: Pectoral fin ^d	RNA ISH; RT-qPCR; skeletal staining	(Leong, Skinner, Shelling, & Love, 2013)

TABLE 1 (Continued)

Syndrome	Human gene	Inheritance pattern	Major human craniofacial/limb phenotypes	Zebrafish ortholog	Model	Zebrafish craniofacial phenotypes	Zebrafish pectoral fin phenotypes	Experimental assays	Reference
Ectrodactyly-ectodermal dysplasia-Clefting syndrome 3 (EEC3) (MIM #604292)	TP63	AD	Ectodermal dysplasia; Orofacial cleft; ectrodactyly	tp63	MO	N/A	Absent: Fin fold; pectoral fin	RT-qPCR; RNA ISH; IHC; BrdU assay; TUNEL assay	(Lee & Kimelman, 2002)
					MO	N/A	Absent: Fin fold; pectoral fin	Transactivation assay; EMSA assay; RNA ISH	(Bakkers, Hild, Kramer, Furutani-Seiki, & Hammerschmidt, 2002)
					LOF CRISPR-Cas9 mutant	N/A	Hypoplastic: Fin fold; Absent: Pectoral fin	RNA-seq; ChIPmentation; ATAC-seq	(Santos-Pereira, Gallardo-Fuentes, Neto, Acemel, & Tena, 2019)
Greig cephalopolysyndactyly syndrome (MIM #175700)	GLI3	AD	Macrocephaly; bulging forehead; broad nasal bridge; wide-set eyes; polydactyly; syndactyly; wide thumb or big toe	gli3	MO	N/A	N/A	RNA ISH; IHC	(Tyrulina et al., 2005)
Hypochondroplasia (MIM #146000)	FGFR3 ^b	AD	Macrocephaly; brachydactyly; bowed legs; shortened limbs	fgfr3	See above.	See above.	See above.	See above.	See above.
Jackson-Weiss syndrome (MIM #123150)	FGFR2 ^a	AD	Craniosynostosis; wide-set eyes; bulging forehead; brachydactyly and/or syndactyly of toes only	fgfr2	See above.	See above.	See above.	See above.	See above.
Kabuki syndrome I (MIM #147920)	KMT2D	AD	Orofacial cleft; microcephaly; micrognathia; wide-set, upward slanting eyes; depressed nose; shortened pinky finger	kmt2d	CRISPR-Cas9 mutant	Hypoplastic/deformed: Ethmoid plate; trabeculae; palatoquadrate	Hypoplastic/deformed: Pectoral fin ^d	RNA-seq; RT-qPCR; IHC; skeletal staining; drug treatment	(Serrano, Demarest, Tone-Pah-Hote, Tristani-Firouzi, & Yost, 2019)
						Absent: Meckel's cartilage; ceratobranchial arches; ceratohyal; hyosymplectic cartilage			
Kabuki syndrome II (MIM #300867)	KDM6A	XLD	Orofacial cleft; microcephaly; micrognathia; wide-set, upward slanting eyes; depressed nose; shortened pinky finger	kdm6a/ kdm6al	MO	Hypoplastic/deformed: Ceratobranchial arches; Meckel's cartilage; ceratohyal (kdm6a only)	Hypoplastic: Pectoral fin (kdm6a only) ^d	Skeletal staining: H&E stain; IHC; live-cell imaging	(Van Laarhoven et al., 2015)
Lacrimalo-dento-digital syndrome (MIM #149730)	FGFR2 ^a	AD	Hypoplasia/aplasia of lacrimal system; orofacial cleft (variable); low-set ears; polydactyly, syndactyly, and/or ectrodactyly	fgfr2	See above.	See above.	See above.	See above.	See above.
	FGFR3 ^b	AD		fgfr3	See above.	See above.	See above.	See above.	See above.

Syndrome	Human gene	Inheritance pattern	Major human craniofacial/limb phenotypes	Zebrafish ortholog	Model	Zebrafish craniofacial phenotypes	Zebrafish pectoral fin phenotypes	Experimental assays	Reference
Muenke syndrome (MIM #602849)	<i>FGFR3</i> ^b	AD	Craniosynostosis; wide-set, bulging eyes; macrocephaly; elevated palate; brachydactyly;	<i>fgfr3</i>	See above.	See above.	See above.	See above.	See above.
Oral-facial-digital syndrome I (MIM #311200)	<i>OFD1</i>	XLD	Orofacial cleft; split tongue; broad nose; wide-set eyes; micrognathia; brachydactyly, syndactyly, clinodactyly	<i>ofd1</i>	MO	Deformed; Meckel's cartilage	N/A	RT-qPCR; RNA ISH; IHC; live-cell imaging; skeletal staining; microarray; bead & dextran injections	(Ferrante et al., 2009)
Pfeiffer syndrome (MIM #101600)	<i>FGFR1</i>	AD	Craniosynostosis; bulging, wide-set eyes; high forehead; micrognathia; beaked nose; brachydactyly; syndactyly	<i>fgfr1a/fgfr1b</i>	MO (<i>fgfr1a</i>) Inducible DN mutant (<i>fgfr1a</i>)	Hypoplastic; Meckel's cartilage Absent: Ceratobranchial arches Hypoplastic/deformed; Meckel's cartilage; ceratohyal; ceratobranchial arches; hyosymplectic cartilage Absent: Meckel's cartilage; ceratohyal; ceratobranchial arches; hyosymplectic cartilage	Hypoplastic; Pectoral fin ^d Hypoplastic; Pectoral fin ^d Absent: Pectoral fin ^d	RNA ISH; RNA FISH; skeletal staining	(Larbuissou et al., 2013)
					CRISPR-Cas9 LOF single mutant	No phenotype	No phenotype	RNA ISH; skeletal staining	(Leerberg et al., 2019)
					CRISPR-Cas9 LOF double mutants (<i>fgfr1a</i> -/-; <i>fgfr1b</i> -/-)	Deformed/hypoplastic; Meckel's cartilage; ceratohyal; hyosymplectic cartilage; ethmoid plate Absent: Ceratobranchial arches	Absent; Pectoral fin		
				<i>fgfr2</i>	See above.	See above.	See above.	See above.	See above.
Popliteal pterygium syndrome (MIM #119500)	<i>IRF6</i> ^c	AD	Orofacial cleft; micrognathia; thin upper lip; syndactyly	<i>irf6</i>	N/A MO DN mRNA OE	N/A No phenotype Hypoplastic; Meckel's cartilage; ceratohyal; palatoquadrate; ceratobranchial arches; hyosymplectic	N/A No phenotype Hypoplastic; Pectoral fin	RNA ISH RT-qPCR; RNA ISH; skeletal staining; TUNEL assay	(Ben, Jabs, & Chong, 2005) (Sabel et al., 2009)
					DN Tol2 mutant	Hypoplastic/deformed; Ethmoid plate; Meckel's cartilage; palatoquadrate	N/A	RNA ISH; IHC; EdU assay; TUNEL assay; live-cell imaging; skeletal staining	(Dougherty et al., 2013)

(Continues)

TABLE 1 (Continued)

Syndrome	Human gene	Inheritance pattern	Major human craniofacial/limb phenotypes	Zebrafish ortholog	Model	Zebrafish craniofacial phenotypes	Zebrafish pectoral fin phenotypes	Experimental assays	Reference
Roberts syndrome (MIM #268300)	ESCO2	AR	Orofacial cleft; micrognathia; downward slanting, wide-set eyes; beaked nose; microcephaly; hypoplastic limbs; syndactyly; ectrodactyly; joint deformities	<i>esco2</i>	MO	Hypoplastic/deformed: Ceratohyal; ceratobranchial arches Absent: Ceratobranchial arches	Hypoplastic/deformed: Pectoral fin	Microarray; RNA ISH; skeletal staining; BrdU assay; TUNEL assay; IHC; RT-qPCR; Caspase assay	(Monnich, Kuriger, Print, & Horsfield, 2011)
Robinow syndrome (MIM #180700)	WNT5A	AD	Prominent forehead; small midface; wide-set, prominent eyes; short nose; brachydactyly	<i>wnt5a</i>	mRNA OE	N/A	N/A	Acridine orange stain; IHC; Western blot; transgene mRNA injections	(Percival et al., 2015)
Saethre-Chotzen syndrome (MIM #101400)	FGFR2 ^a FGFR3 ^b TWIST1	AD AD AD	Craniostenosis; maxillary hypoplasia; high forehead; wide-set eyes; ptosis; broad nasal bridge; brachydactyly; syndactyly; pre-axial polydactyly	<i>fgfr2</i> <i>fgfr3</i> <i>twist1a/twist1b</i>	See above. See above. MO (<i>twist1a</i> ; <i>twist1b</i>)	See above. See above. Absent: Meckel's cartilage; ceratohyal; ceratobranchial arches; palatoquadrate; hyosymplectic cartilage; ethmoid plate; trabeculae	See above. See above. Absent: Pectoral fin ^d	See above. See above. RNA ISH	See above. See above. (Das & Crump, 2012)
Thanatophoric dysplasia I (MIM #187600)	FGFR3 ^b	AD	Macrocephaly; low nasal bridge; brachydactyly; syndactyly; shortened limbs; curved thigh bones	<i>fgfr3</i>	See above.	See above.	See above.	Skeletal staining; live-cell imaging; BrdU assay; RNAscope ISH	(Teng et al., 2018)
Thanatophoric dysplasia II (MIM #187601)	FGFR3 ^b	AD	Cloverleaf skull; macrocephaly; brachydactyly; shortened limbs	<i>fgfr3</i>	See above.	See above.	See above.	See above.	See above.

TABLE 1 (Continued)

Syndrome	Human gene	Inheritance pattern	Major human craniofacial/limb phenotypes	Zebrafish ortholog	Model	Zebrafish craniofacial phenotypes	Zebrafish pectoral fin phenotypes	Experimental assays	Reference
Van der Woude syndrome 1 (MIM #119300)	IRF6 ^c	AD	Orofacial cleft; lower lip pits and/or sinuses; hypodontia; syndactyly	irf6	See above.	See above.	See above.	See above.	See above.
Van der Woude syndrome II (MIM #606713)	GRHL3	AD	Orofacial cleft; lower lip pits and/or sinuses; hypodontia; syndactyly	grhl3	MO	Hypoplastic/deformed: Ceratohyal; ceratobranchial arches; palatoquadrate; Meckel's cartilage; ethmoid plate, Absent: Ceratobranchial arches	Hypoplastic: Pectoral fin ^d	RNA ISH; skeletal staining; electron microscopy; TUNEL assay; micro-ChIP assay; EdU assay;	(Dworkin et al., 2014)
					DN mRNA OE	N/A	N/A	RNA ISH	(Peyrard-Janvid et al., 2014)
					MO; LOF CRISPR-Cas9 mutant	N/A	N/A	IHC	(Miles et al., 2017)

Abbreviations: AD, autosomal dominant; AR, autosomal recessive; CHIP, chromatin immunoprecipitation; DN, dominant negative; EMSA, electrophoresis mobility shift assay; GOF, gain-of-function; H&E, hematoxylin and eosin; IHC, immunohistochemistry; ISH, in situ RNA hybridization; LOF, loss-of-function; MIM, Mendelian Inheritance in Man number; MO, morpholino; N/A, not applicable; OE, overexpression; RNA-FISH, RNA fluorescent in situ hybridization; RT-qPCR, real-time quantitative polymerase chain reaction; TALENs, transcription activator-like effector nucleases; XLD, X-linked dominant.

^aFGFR2 variants are thought to cause nonspecific craniosynostosis and are associated with multiple different craniofacial syndromes.

^bFGFR3 variants are associated with multiple different craniofacial syndromes.

^cIRF6 variants are associated with both popliteal pterygium syndrome and Van der Woude syndrome.

^dOur interpretation of the structure of the pectoral fin based on the Alcian blue skeletal staining images provided in the article.

In this review, we analyze syndromes that present with both craniofacial and limb defects, and we focus specifically on those that have been studied using a zebrafish model. Zebrafish serve as a useful developmental model to study these syndromes because they are relatively easy to genetically modify, inexpensive to maintain, reproduce and develop quickly, and the transparency of the embryo allows for live imaging *in vivo*. The zebrafish skeleton is also simpler, with fewer structural elements and larval cartilages are only a few cell layers thick, making them easier to visualize. Its pectoral and pelvic fins are homologous to mammalian forelimbs and hindlimbs, respectively (Grandel & Schulte-Merker, 1998). Most of the gene regulatory networks involved in both craniofacial and limb/fin development are highly conserved between zebrafish and mammals. The information obtained from zebrafish studies can provide valuable insight into human development. First, we will very briefly describe the processes governing craniofacial and limb/fin development as a preface. Then we will highlight recent studies that have used zebrafish to uncover the mechanism(s) by which certain genes cause syndromes with craniofacial and limb defects. We hope to identify common themes and mechanisms that can help explain why these two defects often occur together.

2 | BRIEF OVERVIEW OF CRANIOFACIAL AND LIMB DEVELOPMENT

2.1 | Craniofacial development

Much of the facial skeleton is derived from cranial neural crest cells. Neural crest cells (NCCs) are a multipotent population of cells that originate at the dorsal most part of the neural tube. During development, NCCs undergo an epithelial-mesenchymal transition to delaminate from the neural tube, migrate to different regions of the body, and differentiate into specific cell types. Cranial NCCs are a subpopulation of NCCs that migrate from the neural tube into the pharyngeal arches and the facial prominences. They interact with head mesoderm in response to signals from the ectoderm and endoderm to differentiate into cartilage, bone, cranial neurons, glia, and connective tissues to form the frontonasal skeleton, jaw, odontoblasts of the teeth, middle ear, glia, and cranial neurons.

The skull is divided into two structures, the viscerocranium and neurocranium. The viscerocranium forms the lower part of the skull and supports the structure of the face. It is comprised entirely of NCCs from the pharyngeal arches (Kague et al., 2012; Morriss-Kay, 2001). Each pharyngeal arch gives rise to different structures. The first arch forms the lower jaw (mandibular domain) and the palate (maxillary domain); the second arch (hyoid) forms the ceratohyal and hyomandibular bones, which connect the lower jaw to the neurocranium; and in zebrafish, the third through the seventh arches give rise to the ceratobranchial cartilages, which support the gill tissues (Schilling & Kimmel, 1997) (Figure 1a,c). In mammals, these posterior arches form laryngeal cartilage. Interestingly, parts of the zebrafish jaw and hyoid skeleton are evolutionarily homologous to the mammalian middle ear bones. More specifically, the Meckel's cartilage and





FIGURE 1 Homologous craniofacial structures between zebrafish and mammals. (a) Lateral view of zebrafish head skeleton. The posterior end of the Meckel's cartilage, the palatoquadrate and hyosymplectic cartilage are homologous to structures in the mammalian middle ear, particularly the malleus, incus, and stapes bone (b). (c) Dorsal view of the zebrafish neurocranium, which is analogous to the mammalian palate (d). (e,f) Dorsal view of the zebrafish and mammalian cranial sutures. Homologous structures have the same color, and analogous structures have added hashmarks. Adapted from Mork and Crump (2015). c, cochlea; cbs, ceratobranchial arches #1–5; ch, ceratohyal; cs, coronal sutures; ed, ear drum; ep, ethmoid plate; fr, frontal bones; hs, hyosymplectic cartilage; i, incus; ifs, interfrontal sutures; ls, lambdoid sutures; m, Meckel's cartilage; ma, malleus; mep, medial ethmoid plate; n, notochord; oc, occipital bone; pa, parietal bones; pch, parachordal; pp, primary palate; pq, palatoquadrate; ps, parasphenoid; so, supraoccipital bone; sp, secondary palate; ss, sagittal sutures; st, stapes bone; tr, trabecula

part of the palatoquadrate are homologous to the malleus and incus of the middle ear, while the hyomandibula portion of the hyosymplectic cartilage is homologous to the stapes bone (Figure 1a, b) [reviewed in Anthwal, Joshi, & Tucker, 2012, Mork & Crump, 2015]. Molecular mechanisms governing lower jaw development have been well-studied in both mice and zebrafish. For a more detailed review, we refer to Clouthier, Garcia, and Schilling (2010) and Santagati and Rijli (2003). One of the most critical factors in lower jaw development is endothelin-1 (*Edn1*) signaling through the endothelin-A receptor (*Ednra*) (schematic in Figure 2a). In zebrafish, loss of *edn1* leads to a severe truncation of the Meckel's cartilage, a loss of the ceratohyal bone, and a homeotic transformation of the

lower jaw to an upper jaw (Kimmel, Ullmann, Walker, Miller, & Crump, 2003; Miller, Schilling, Lee, Parker, & Kimmel, 2000). Similar phenotypes are observed in *Edn1*^{−/−} and *Ednra*^{−/−} mice, suggesting that their roles in craniofacial development are conserved (Clouthier et al., 1998; Kurihara et al., 1994). Endothelin signaling is required for downstream expression of *Dlx5/Dlx6*, as well as *Dlx3* and *Hand2* in the ventral most part of the pharyngeal arch. The *dlx* homeobox genes are important for establishing the dorsal/ventral axis in craniofacial development and are further regulated by *Mef2c* (Miller et al., 2007). *nkx3.2* (*bapx1*) is an additional homeobox gene that regulates genes involved in jaw joint formation. Its expression is limited to the intermediate region of the pharyngeal arch, and it is repressed by *Hand2* in

FIGURE 2 Molecular mechanisms governing lower jaw and cranial suture development. (a) Simplified schematic of endothelin-1 signaling in the first pharyngeal arch during lower jaw development. Endothelin signaling is required for downstream expression of *Dlx* homeobox genes, which are required for dorsal/ventral patterning, and *Hand2*, which represses *Nkx3.2* in the ventral region of the pharyngeal arch. *Nkx3.2* is also regulated by *Bmp4* in the most dorsal part of the arch and *Fgf8* in the ectoderm. (b) Schematic of genes and pathways involved in cranial suture development. *Twist1* is capable of forming homodimers (T/T) and heterodimers with E proteins (T/E). These two forms act antagonistically with one another to regulate *FGFR2* activity, downstream BMP signaling, osteogenic differentiation, and suture closure. ID, inhibitor of DNA binding; ph, pharyngeal arch; T/E, *Twist1* heterodimer; T/T, *Twist1* homodimer



the ventral region, *Fgf8* in the oral epithelium, and *Bmp4* in the distal pharyngeal arch (Miller, Yelon, Stainier, & Kimmel, 2003; Miyashita et al., 2020; Wilson & Tucker, 2004). *DLX5/DLX6* and *NKX3.2* are also involved in human limb development. Variants are associated with split hand/foot malformation (MIM #183600) and spondylo-megaepiphyseal-metaphyseal dysplasia (MIM #613330), respectively.

The neurocranium is composed of both cranial NCCs and mesoderm and protects and supports the brain (Wada et al., 2005). Cranial NCCs in the maxillary domain of the first pharyngeal arch give rise to the palate after cells migrate medially to converge at the midline of the roof of the mouth. In mammals, palatal shelves composed of cranial NCCs grow and converge at the midline to form an epithelial seam, while the primary and secondary palates fuse to create the palatal skeleton (Figure 1d). The zebrafish palate is located in the anterior part of the neurocranium and consists of the ethmoid plate, trabeculae, and parasphenoid bone (Figure 1c). Orofacial clefting occurs when the palatal shelves fail to come together. Clefting, truncation,

hypoplasia, or the absence of these structures is indicative of orofacial clefting in zebrafish (Dougherty et al., 2013; Eberhart et al., 2008; Swartz, Sheehan-Rooney, Dixon, & Eberhart, 2011; Wada et al., 2005). It is still unclear whether cranial NCCs that form the palate fuse in zebrafish as well or if they are differentiating in a posterior to anterior pattern (Dougherty et al., 2013; Swartz et al., 2011). Early palatogenesis is conserved among vertebrates. Mice and zebrafish share similar expression patterns of critical genes involved in palate formation in the anterior maxillary domain (*msxe*, *bmp4*, *bmp2b*, and *fgf10a*) as well as the posterior maxillary domain (*tbx22*, *osr1*, *osr2*, *pax9a*) (Braybrook et al., 2002; Peters, Neubuser, Kratochwil, & Balling, 1998; Swartz et al., 2011) [reviewed in (Hilliard, Yu, Gu, Zhang, & Chen, 2005)]. They also utilize the same signaling pathways, such as *Fgf*, *Pdgr*, *Bmp*, *Tgfb*, *Wnt*, and *Shh*, and disruptions to any of these pathways result in craniofacial defects, particularly in the anterior neurocranium (Dougherty et al., 2013; Eberhart et al., 2008; Swartz et al., 2011; Wada et al., 2005). For example, *fgf10a* is

expressed throughout the oral ectoderm and knocking down this gene produces a shortened trabeculae and parasphenoid bone and a misshapen Meckel's cartilage and palatoquadrate (Swartz et al., 2011). Fgf10a regulates Shh signaling, which is critical for cranial NCC migration to the midline and induction of chondrogenesis. Loss of Shh in zebrafish causes inappropriate fusion of the trabeculae (Wada et al., 2005). Interestingly, Fgf10a and Shh are both critical for limb development as well (see below). Loss of either gene leads to defects in the pectoral fin, which is homologous to mammalian forelimbs (Swartz et al., 2011; van Eeden et al., 1996). Although early development of the mammalian and zebrafish palates appear genetically similar with similar patterning formation, there are distinct morphogenic differences at later stages, warranting caution when comparing zebrafish and mammalian palatogenesis [reviewed in (Bush & Jiang, 2012)].

The skull vault of the neurocranium has five bones that are connected by cranial sutures, or fibrous tissues. The anatomical structure of the skull vault and its cranial sutures is conserved between humans and zebrafish (Figure 1e,f). The sutures are patent in early development, providing room for the skull and brain to grow. The timing of suture closure differs as human sutures are only patent during early childhood whereas zebrafish sutures remain patent throughout the life of the fish (Quarto & Longaker, 2005). The molecular mechanisms dictating cranial suture formation and closure are conserved among vertebrates, rendering zebrafish a useful model for studying craniosynostosis, a common skeletal defect that occurs when cranial sutures prematurely fuse (Quarto & Longaker, 2005; Topczewska, Shoela, Tomaszewski, Mirmira, & Gosain, 2016) [reviewed in (Holmes, 2012)]. The skull stops growing perpendicular to the fused suture and compensates by growing in a parallel fashion. Therefore, patients with craniosynostosis have abnormally shaped skulls. Most cases of craniosynostosis are caused by genetic variants in *TWIST1*, *FGFRs*, and *EFNB1* among other genes [reviewed in Wu & Gu, 2019]. Twist1 is a transcription factor that is expressed in either the osteogenic front as a homodimer (T/T) or as a heterodimer with E proteins (T/E), such as Tcf12, in the mesenchymal cells of cranial sutures (schematic in Figure 2b). These two forms of Twist1 act in opposition to one another. The T/T homodimers activate *FGFR2* and promote osteogenic differentiation by increasing BMP signaling as well as *Msx2* and *Runx2* expression. This leads to suture closure. In contrast, the T/E heterodimer represses *FGFR2*, preventing osteogenesis and suture closure. The ratio of Twist1 homodimer to heterodimer changes over time as an organism grows and develops. An untimely excess of T/T homodimer leads to increased *FGFR2* expression, decreased Eph/Ephrin signaling, an inappropriate influx of neural crest cells to the paraxial mesoderm, and craniosynostosis (Connerney et al., 2008; Merrill et al., 2006).

Later in this review, we summarize studies that have used zebrafish to study syndromes with craniofacial anomalies, including orofacial clefting and craniosynostosis.

2.2 | Limb development


The zebrafish pectoral and pelvic fins are homologous to mammalian forelimbs and hindlimbs, respectively. Limb growth begins at the

lateral plate mesoderm, where mesenchyme precursors form a small bud surrounded by an ectodermal layer (schematic in Figure 3d). Retinoic acid is synthesized in the surrounding somites and, along with Wnt signaling, establishes the limb field and initiates limb induction (Grandel et al., 2002; Ng et al., 2002). Fgf signaling is required for the formation and function of the apical ectodermal ridge (AER) (Niswander, Tickle, Vogel, Booth & Martin 1993; Sekine et al., 1999; Sun, Mariani, & Martin, 2002), which regulates limb outgrowth and establishes the proximodistal axis (shoulder to digits) (Saunders, 1948). Transcription factors Tbx5 (Agarwal et al., 2003; Ng et al., 2002) and p63 (Bakkers et al., 2002) induce Fgf signaling in the mesenchyme and AER. This establishes a complex epithelial-mesenchymal feedback loop that activates proliferation and differentiation of mesenchymal cells resulting in limb outgrowth (Ohuchi et al., 1997). Fgf signaling in the AER is negatively regulated by BMP signaling (Niswander & Martin, 1993; Pajni-Underwood, Wilson, Elder, Mishina, & Lewandoski, 2007; Pizette & Niswander, 1999). Moreover, BMP signaling is required for interdigital programmed cell death and preventing finger webbing as well as polydactyly (Pajni-Underwood et al., 2007; Selever, Liu, Lu, Behringer, & Martin, 2004). The anterior/posterior axis (digits 1–5) is established by Shh signaling in the zone of polarizing activity (ZPA) (Riddle, Johnson, Laufer, & Tabin, 1993; Saunders & Gasseling, 1968). Shh also regulates and is regulated by Fgf signaling in the AER (Laufer, Nelson, Johnson, Morgan, & Tabin, 1994; Niswander, Jeffrey, Martin, & Tickle, 1994). The dorsal/ventral axis (back of hand to palm) is dictated by expression of Wnt7a (dorsal) and En1(ventral) (Davis, Holmyard, Millen, & Joyner, 1991; Gardner & Barald, 1992; Loomis, Kimmel, Tong, Michaud, & Joyner, 1998; Parr & McMahon, 1995). Each gene and pathway are interconnected, and dysregulation at any point can cause abnormal limb growth [reviewed in (Kantaputra & Carlson, 2019)] (Figure 3d).

The first skeletal elements to form in the zebrafish pectoral fin bud are the scapulocoracoid, postcoracoid, and endoskeletal disk, which are all derived from fin mesenchymal cells. By 4 dpf (days post fertilization), collagenous rays called actinotrichia form and act as supportive elements for the fin fold (Figure 3a). This larval endoskeleton structure persists for several weeks until cells in the endoskeletal disk proliferate and expand the fin. The cartilage matrix of the intermediate larval structure decomposes, and at 23 dpf (7.1 mm standard length), rod-like dermal bones called lepidotrichia, or fin rays, form. By 28 dpf (9.5 mm), the endoskeletal disk splits to form four proximal radial bones. There are also six to eight distal radial bones located above and distal to the proximal radials that later fuse to the lepidotrichia (Grandel & Schulte-Merker, 1998) (Figure 3b). The pelvic fins do not begin formation until 18 dpf (6 mm), and their development is similar to that of pectoral fins. The main difference is that the pelvic fin bud develops more quickly and does not require an intermediate larval endoskeleton before the terminal adult fin structure (29 dpf, 10.1 mm). For a more detailed description of the anatomy of zebrafish fin development, we refer to Grandel and Schulte-Merker (1998).

Morphologically, zebrafish fins and mammalian limbs are distinct from one another, and it is difficult to assign which elements are truly

FIGURE 3 The gene regulatory networks involved in zebrafish and mammalian limb development are highly conserved. The zebrafish pectoral fin, shown at 4 days post fertilization (a) and adulthood (b), is homologous to the mammalian forelimb (c). (d) Schematic of genes and pathways involved in early limb development. Limb bud initiation begins with retinoic acid (RA) signaling in the surrounding somites (teal). This then leads to the initiation of Fgf signaling in the mesoderm and ectoderm and limb bud growth at the lateral plate mesoderm. The AER (navy) dictates outgrowth and the proximal/distal axis; the ZPA (purple) determines the anterior/posterior axis; and WNT7A/EN1 (green) establish the dorsal/ventral axis. As such, each gene and pathway are interconnected and rely on one another for proper limb growth. Abbreviations: ac, actinotrichia; AER, apical ectodermal ridge; c, carpal bones; dr, distal radials; ed, endoskeletal disk; fr, fin rays; h, humerus; mc, metacarpals; pc, postcoracoid; ph, phalanges; pr, proximal radials; r, radius; RA, retinoic acid; sc, scapulocoracoid; u, ulna; ZPA, zone of polarizing activity



homologous [reviewed in (Yano & Tamura, 2013)] (Figure 3a–c). Nevertheless, the conservation of the gene regulatory networks indicates that they are still a useful model for limb development. Researchers can examine the larval pectoral fin (4–5 dpf) and look for deformities in the shape and length of cartilage structures. If a mutant survives to adulthood, researchers can look more closely at the bones that develop later and use them as a model for human limb bones. Differences in the number of proximal radial bones, number of distal radial bones, or the length of lepidotrichia can be indicators of human polydactyly (more bones), ectrodactyly (fewer bones), or brachydactyly (shortening of bones). Medaka fish (*Oryzias latipes*) has been used by Letelier et al. (2018) to model human ectrodactyly caused by decreased *SHH* signaling by partially deleting an upstream *shh* limb enhancer known as ZRS. This deletion leads to a decrease in proximal radial bones from four to two in adult fish. Below, we discuss ways in

which zebrafish have been used to study human craniofacial and limb defects.

3 | ZEBRAFISH MODELS OF CRANIOFACIAL ANOMALIES WITH ACCOMPANYING LIMB DEFECTS

Structures of the craniofacial and limb skeleton are clearly distinct from one another, but they utilize many of the same gene regulatory networks and mechanisms for their development, such as Fgf, Bmp, Wnt, and Shh signaling. It follows then that several human congenital syndromes caused by single gene variants present with defects in both structures (Panthaki & Armstrong, 2003). Here we will describe a few examples in which zebrafish were used to study specific

syndromes characterized by orofacial clefting, craniosynostosis, and limb defects. Orofacial clefting and craniosynostosis are the two most common craniofacial phenotypes, so our primary focus is here. A full list of craniofacial and limb human syndromes that have been modeled with zebrafish can be found in Table 1. Some of these studies explored different phenotypes associated with the syndrome. For example, Fuller et al. (2014) use zebrafish to model the left/right patterning defects observed in patients with Carpenter syndrome I (MIM #201000). We have chosen to highlight studies that specifically studied genes in the context of craniofacial and/or limb development.

3.1 | Orofacial clefting and missing/fused digits

Orofacial clefting, namely cleft lip with or without cleft palate, is the most common craniofacial anomaly occurring in 1 in 700 newborn babies (Global registry and database on craniofacial anomalies, 2001). Cleft palate occurs in humans when the palatal shelves fail to elevate or properly fuse at the midline. As mentioned above, the zebrafish ethmoid plate is often used to model the mammalian palate. For a comprehensive review on zebrafish models of orofacial clefting we refer to Duncan, Mukherjee, Cornell, and Liao (2017). Here, we will discuss recent zebrafish studies of syndromes that present with both orofacial clefting and limb defects, primarily ectrodactyly (missing digits) and syndactyly (fused digits).

3.1.1 | Roberts syndrome

Roberts syndrome (MIM #268300) is an autosomal recessive disorder characterized by orofacial clefting, micrognathia (hypoplastic jaw), downward slanting, wide-set eyes, a beaked nose, microcephaly, hypoplastic limbs, syndactyly, and joint deformities (Roberts, 1919). Using multipoint linkage analysis, Vega et al. (2005) found eight different homozygous variants in *ESCO2* in affected individuals. *ESCO2* is a highly conserved gene that encodes an N-acetyltransferase required for holding sister chromatids together after DNA replication and before mitosis (Rolef Ben-Shahar et al., 2008). In zebrafish, *esco2* is expressed in the branchial arches and pectoral fins, as well as brain ventricles, optic vesicles and retinal cells during early development (Monnich et al., 2011). Partial knockdown of *esco2* in zebrafish morphants (organisms treated with morpholino antisense oligonucleotides to knockdown gene expression) leads to disorganized craniofacial cartilage and hypoplastic jaw elements as well as shortened, abnormally shaped pectoral fins. This is due to cells being blocked at the onset of mitosis, leading to increased cell death. *esco2* morphant cells in mitosis have distorted, disorganized mitotic spindles, likely from the instability of the sister chromatids (Vega et al., 2005). Consistent with this study, stable *esco2*^{-/-} mutants created with CRISPR-Cas9 also have smaller heads, missing pectoral fins, and increased apoptosis, particularly at the neural tube (Percival et al., 2015). This group observed that cells are trapped in mitosis. Indeed, after the nuclear envelope breaks down, chromosomes in *esco2*^{-/-} mutants

scatter and are not captured on the metaphase plate, blocking mitosis from proceeding. This leads to aneuploidy and/or micronuclei and the cells are forced to undergo apoptosis. The craniofacial and limb defects, particularly the hypoplastic jaw and limbs, observed in patients from loss of *ESCO2* are hypothesized to result from increased cell death, based on these zebrafish studies.

3.1.2 | Van der Woude syndrome

Van der Woude syndrome is an autosomal dominant disorder caused by variants in either *IRF6* (MIM #119300) (Burdick, Bixler, & Puckett, 1985; Kondo et al., 2002) or *GRHL3* (MIM #606713) (Peyrard-Janvid et al., 2014). Patients present with orofacial clefting, lower lip pits and/or sinuses, hypodontia, and syndactyly.

IRF6 is a member of the interferon regulatory transcription factor family. In zebrafish larvae, it is expressed in the pharyngeal arches and in the epithelial cells of the mouth, esophagus, and pharynx, as well as the olfactory and otic placodes (2–72 hpf) (Ben et al., 2005). Expression of *irf6* in the pectoral fin has not yet been shown; however, injection of a dominant-negative form of *irf6* RNA into zebrafish embryos at the single-cell stage leads to shortened/loss of the pectoral fin as well as hypoplastic, disorganized craniofacial elements and a clefted ethmoid plate (Sabel et al., 2009). Dougherty et al. (2013) created stable, dominant-negative *irf6* mutants driven by a *sox10* promoter to limit mutant expression to NCCs. With time-lapse imaging, they show that chondrocytes at the median and lateral ethmoid plate fail to come together in mutants, creating a cleft (Dougherty et al., 2013). Similarly, *Irif6*^{-/-} mutant mice have clefting in the secondary palate and a hypoplastic snout and jaw (Ingraham et al., 2006). *Irif6* is thought to be involved in endothelin signaling during palate formation (Fakhouri et al., 2017). The clefting phenotypes observed in zebrafish and mice are consistent with what is seen in Van der Woude patients and demonstrate an important role for *IRF6* in palatogenesis.

grhl3 is a transcription factor selectively expressed in the non-neural ectoderm as well as endoderm pouches surrounding the pharynx of developing zebrafish embryos. Knockdown of *grhl3* causes severe hypoplasia of the palatoquadrate, ceratohyal, and Meckel's cartilage and loss of the ceratobranchial arches and pectoral fins (Dworkin et al., 2014). There is increased cell apoptosis in the pharyngeal arches with loss of *grhl3*. Using a micro-ChIP assay, Dworkin et al. (2014) found that Grhl3 directly binds to the promoter of *edn1*, a highly conserved gene required for lower jaw development (Clouthier et al., 1998; Miller et al., 2000). Expression of *edn1* and its known downstream targets, *hand2* and *dlx3*, are significantly reduced in the endoderm of *grhl3* morphants. Importantly, injection of *edn1* mRNA into *grhl3* morphants rescues *hand2* and *dlx3* expression as well as the craniofacial and limb skeletons (Dworkin et al., 2014). It is thought that *grhl3* is required for *edn1* expression in the pharyngeal endoderm, which is then important for NCC growth and proliferation and palatogenesis.

irf6 and *grhl3* are likely required for limb development in zebrafish, as shown by the loss of pectoral fins in different models,

but their roles remain unclear. During mammalian development, loss of either *Irf6* or *Grlh3* results in shortened forelimbs, ectrodactyly, or syndactyly (Ingraham et al., 2006; Kashgari et al., 2020). *Irf6* is thought to be required for formation of the periderm, a single layer of epithelial cells surrounding the embryonic epidermis. In mice, *Irf6* directly regulates transcription of *Grlh3* (de la Garza et al., 2013), which was recently shown to be required for digit separation (Kashgari et al., 2020). It is unknown whether the interaction between *irf6* and *grhl3* during limb formation is conserved and if this accounts for the loss of pectoral fins seen in zebrafish.

3.1.3 | Ectrodactyly-ectodermal dysplasia-clefting syndrome 3 (EEC3)

Ectrodactyly-ectodermal dysplasia-clefting syndrome 3 (EEC3) (MIM #604292) is an autosomal dominant disorder caused by mutations in *TP63*, a transcription factor and member of the *TP53* family (Celli et al., 1999). EEC3 is characterized by orofacial clefting, ectrodactyly, and ectodermal dysplasia such as hypopigmented, scaly skin and malformed/decayed teeth (Penchaszadeh & de Negrotti, 1976). *TP63* is transcribed from two different promoters, creating two isoforms of the gene, *Tap63* and Δ Np63, which then act as an activator or repressor, respectively (Yang et al., 1998). Zebrafish studies have shown that knockdown of the dominant negative repressive isoform, Δ Np63, results in loss of the pectoral fin and a decrease in proliferation in the epidermis (Bakkers et al., 2002; Lee & Kimelman, 2002). p63 loss-of-function mutants (*tp63*^{-/-}) created with CRISPR-Cas9 recapitulate this phenotype (Santos-Pereira et al., 2019). Unfortunately, the authors did not note any craniofacial defects in the *tp63*^{-/-} mutants, and this is likely due to the larvae dying between 40–50 hours post fertilization (hpf). *tp63*^{-/-} mutants have a significant decrease in expression of epidermal genes as well as genes relating to fin development. Additionally, Gene Ontology (GO) term enrichment analyses show a downregulation in genes relating to cell-matrix adhesion, cell adhesion, and skeletal muscle fiber development, genes which are likely to contribute to craniofacial and limb development. However, further studies exploring these functions have not yet been explored. *TP63*^{-/-} mutant mice display severe craniofacial and limb defects, including cleft lip and palate, hypoplastic upper and lower jaw, and limb truncations (Yang et al., 1999). Moreover, it has been shown in mice that Δ Np63 directly regulates the transcription of *Dlx5/Dlx6* (Lo Iacono et al., 2008) and *Fgf8/Fgf4* (Kawata et al., 2017), which are all important for both limb and craniofacial development (Figures 2a and 3d). Therefore, it may be worth revisiting the zebrafish model to determine whether craniofacial defects are present in *tp63*^{-/-} mutants and if *TP63* is regulating these key developmental genes.

3.1.4 | Craniosynostosis and variable limb defects

Craniosynostosis is the second most common craniofacial anomaly with a prevalence of 1 in 2,500 live births (Boulet, Rasmussen, &

Honein, 2008). Craniosynostosis is the premature fusion of skull bones at the sutures, or fibrous structures that join the bones, before the brain has fully formed. As a result, the skull is misshapen. During normal development, the sutures hold the skull in place while remaining flexible to allow proper brain growth. The anatomical structure and molecular mechanisms dictating formation of cranial sutures is conserved between humans and zebrafish, making them a useful model for studying craniosynostosis (Quarto & Longaker, 2005). Here, we will discuss Saethre-Chotzen syndrome and syndromes associated with mutations in *FGFR2*.

3.1.5 | Saethre-Chotzen syndrome

Saethre-Chotzen syndrome is an autosomal dominant disorder featuring craniosynostosis, due to loss of the coronal suture, maxillary hypoplasia, a high forehead, wide-set eyes, ptosis (droopy eyelids), a broad nasal bridge, brachydactyly, syndactyly, and polydactyly in the feet (Chotzen, 1932; Saethre, 1931). It is associated with variants in *FGFR2* (see below), *FGFR3*, and *TWIST1* (MIM #101400) (el Ghouzzi et al., 1997; Howard et al., 1997; Paznekas et al., 1998). Transcription factors *Twist1a* and *Twist1b*, the zebrafish orthologs of *TWIST1*, are both required for ectomesenchyme specification from NCCs (Das & Crump, 2012). Knockdown of either gene with morpholinos results in minor skeletal defects, such as minor hypoplasia of the ventral mandibular and hyoid cartilages, but a double knockdown leads to an almost complete loss of the viscerocranium and loss of pectoral fins (Das & Crump, 2012). Stable double *twist1a*^{-/-};*twist1b*^{-/-} knockout mutants created using TALENs (transcription activator-like effector nucleases) have a milder phenotype than the morphants and present with hypoplasia in the Meckel's cartilage, palatoquadrate, and hyosymplectic cartilage (Teng et al., 2018). Using transgenic models and the GAL4:UAS system, Das and Crump (2012) show that *Twist1* promotes ectomesenchyme cell fates (i.e., cartilage, bones, etc.) in cranial NCCs. These cells primarily make up the craniofacial skeleton.

One of the most distinct phenotypes of Saethre-Chotzen syndrome is the selective loss of the coronal suture, which separates the two parietal bones from the frontal bone in the skull. Teng et al. (2018) created viable, triple *tcf12*^{-/-};*twist1a*^{-/-};*twist1b*^{-/-} mutants that show mild craniofacial phenotypes at 5 dpf but adult mutants develop severe unilateral or bilateral coronal synostosis. Juvenile *tcf12*^{-/-};*twist1b*^{-/-} double mutants assessed for mineralization (Calcein green staining), bone (Alizarin red staining), and live-cell imaging of osteoblasts (*sp7:EGFP* transgenic line) all show accelerated growth of the frontal and parietal bones in mutants compared to wildtype. These bones grow diagonally toward one another, becoming aberrantly shaped and leading to premature fusion, which is consistent with the human phenotype. Premature fusion of the suture prevents bone enlargement and growth of the skull along the anterior/posterior axis. Finally, using RNAscope in situ hybridization, a recently developed technique that uses probes to amplify target RNA in intact cells and tissues, they show that *tcf12*^{-/-};*twist1b*^{-/-} mutants have reduced expression of skeletal stem cell markers, *gli1*, *grem1a*, and

prx1a in the coronal sutures as well as a decrease in osteoprogenitors compared to the wildtype. This is true only in the coronal sutures, suggesting that there is a selective exhaustion of osteoprogenitors in this region and bone growth has ceased in the mutants (Teng et al., 2018). These studies demonstrate the importance of *twist1* in craniofacial development.

twist1 also appears to be required for pectoral fin development, as suggested by the skeletal preparations for *twist1a* and *twist1b* morphants (Das & Crump, 2012). However, the mechanism is not well understood. In mice, loss of Twist1 blocks forelimb growth. It is required for AER and ZPA maintenance by regulating Fgf and Shh signaling, respectively (O'Rourke, Soo, Behringer, Hui, & Tam, 2002). Further studies in zebrafish are needed to determine if these interactions are conserved in the growing pectoral fin bud.

3.1.6 | FGFR2 syndromes

It is well-known that fibroblast growth factor (FGF) signaling is critical in many aspects of development. Humans have four tyrosine kinase FGF receptors (FGFR) that can bind to eighteen different secreted FGF proteins. Variants in *FGFR2* specifically are known to be associated with several different craniofacial and limb syndromes, including Apert syndrome (MIM #101200), bent bone dysplasia (MIM #614592), Jackson Weiss syndrome (MIM #123150), Pfeiffer syndrome (MIM #101600), and Saethre-Chotzen syndrome (MIM #101400) [reviewed in Wenger, Miller, & Evans, 1998–(2020)]. Interestingly, each of these syndromes presents with craniosynostosis and either brachydactyly or syndactyly. A more detailed description of phenotypes for each syndrome can be found in Table 1. Mutations in other FGFRs and secreted FGF proteins are also known to lead to developmental disorders [reviewed in Wenger et al., 1998–(2020)].

In zebrafish embryos, *fgfr2* is expressed in the pharyngeal endoderm and hindbrain between 24–72 hpf (Larbuissou et al., 2013) and in the cranial sutures at 6 weeks post fertilization (Topczewska et al., 2016). Treatment of embryos between 18–24 hpf with SU5402, an inhibitor of FGFRs, leads to loss of the pharyngeal arches (Walshe & Mason, 2003). Injection of morpholinos against *fgfr2* leads to hypoplastic skeletal structures, including the Meckel's cartilage and palatoquadrate, as well as shortened pectoral fins. Additionally, knockdown of both *fgfr1a* and *fgfr2* produces an even more severe phenotype with loss of the ceratohyal, ceratobranchial arches, neurocranium, and pectoral fins (Larbuissou et al., 2013). In situ RNA hybridization experiments in single morphants show no changes in expression of *sox9*, *dlx2a*, or *fli1*, suggesting that cranial NCC migration into the pharyngeal arches is not affected by loss of *fgfr2*. However, there is decreased expression of *barx1* and *runx2b*, suggesting Fgfr2 required for chondrocyte condensation and maturation (Larbuissou et al., 2013).

Leerberg et al. (2019) generated knockout zebrafish mutants for single *fgfr* genes using CRISPR-Cas9. Surprisingly, single homozygous mutants are viable and mRNA expression levels of non-mutated *fgfr* genes are not elevated, suggesting genetic redundancy. *fgfr1a*–/–;

fgfr1b–/–;*fgfr2*–/– triple mutants had the most severe phenotypes in the limb and craniofacial skeleton. Consistent with the morphant studies, these mutants had shortened or loss of pectoral fins. By in situ RNA hybridization, there was a significant decrease in *fgf24*, *fgf8a*, and *dlx2a*, which are markers of limb outgrowth. The triple mutants also exhibited severe craniofacial phenotypes, including a loss of the ceratobranchial arches, ceratohyal, and hyosymplectic cartilage; a misshapen palatoquadrate; a downward facing Meckel's cartilage; and a clefted ethmoid plate. This is likely not due to issues with cranial NCC migration, but rather maintenance of the NCCs (Leerberg et al., 2019).

Fgf signaling is critical in both craniofacial and limb development. As shown in these studies and in mouse studies, Fgf signaling is required in the facial skeleton for early patterning, growth regulation, tooth growth, palatogenesis, and suture formation [reviewed in Nie, Luukko, and Kettunen (2006)]. In the limb, it is necessary for proximodistal outgrowth and regulating the anterior/posterior axis [reviewed in Mercader (2007)]. Therefore, mutations in *FGFR* genes can lead to syndromes that present with defects in either or both craniofacial and limb structures.

4 | SUMMARY

In this review, we have provided a brief overview of both craniofacial and limb development and discussed the relevance of using zebrafish as a model for studying both developmental mechanisms. We then reviewed studies that have used zebrafish to study human syndromes that present with both craniofacial and limb defects and attempted to understand why these defects often occur together. We speculate that there is a pleiotropic effect, in which a single gene affects more than one developmental process. Indeed, we have discussed how many signaling pathways, such as Fgf, Shh, Bmp, and Wnt signaling, are involved in both craniofacial and limb development. Disruptions to components of these pathways results in both craniofacial and limb defects. Moreover, some transcription factors, such as Twist1, Irf6, and Tp63 have been shown in mice to bind to and regulate genes that are critical for craniofacial and limb development. Additional studies are required to show that this is conserved in zebrafish as well. NCCs and their derivatives, namely melanoblasts and neurons, can be found in the limb mesenchyme, which may contribute to the similar gene regulatory networks (Erickson, 1985; Grim & Christ, 1993). It has also been proposed that there is a deep homology between the craniofacial and limb skeletons. In cartilaginous fishes (Chondrichthyes), appendages grow out of the gill arches, a structure that later gives rise to part of the craniofacial skeleton. This led Carl Gegenbaur in 1878 to hypothesize that paired limbs/fins are evolved from the gill arches (Gegenbaur, 1878). This idea was recently supported by the work of Gillis and colleagues. In the little skate (*L. erinacea*), they demonstrated that chondrichthyan branchial rays and appendages both rely on a complex interplay between retinoic acid, *Shh* and *Fgf8* signaling to drive endoskeleton outgrowth and patterning of both the gill arches and appendages (Gillis, Dahn, & Shubin, 2009; Gillis & Hall, 2016).

They argue that this is due to a deep homology of the structures. More recently, they have shown in the little skate that the first gill arch is composed of NCCs and the fins are made of mesoderm cells, as expected. Interestingly, the more posterior gill arches are composed of both NCCs and lateral mesoderm cells, suggesting that gills and fins develop from a common pool of cells that have the potential to develop into either structure (Sleight & Gillis, 2020). This may explain why craniofacial and limb patterning are so similar. Despite the seemingly stark differences between humans and zebrafish, we have shown that zebrafish have emerged as a powerful tool to study human craniofacial and limb development.

ACKNOWLEDGMENTS

This work is supported by the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) (1F31HD103368-01 to B.T.T. and 1R03HD096320-01A1 to K.B.A.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. We thank members of the K.B.A. laboratory, particularly Lomeli C. Shull and Ezra S. Lencer; Jennyfer M. Mitchell; and Lee Niswander for critically reviewing the manuscript. We apologize to researchers whose work we were unable to include due to space limitations.

CONFLICT OF INTEREST

The authors declare no potential conflict of interest.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were generated in this study.

ORCID

Kristin B. Artinger  <https://orcid.org/0000-0002-3003-6042>

REFERENCES

Agarwal, P., Wylie, J. N., Galceran, J., Arkhitko, O., Li, C., Deng, C., ... Bruneau, B. G. (2003). Tbx5 is essential for forelimb bud initiation following patterning of the limb field in the mouse embryo. *Development*, 130(3), 623–633. <https://doi.org/10.1242/dev.00191>

Anthwal, N., Joshi, L., & Tucker, A. S. (2012). Evolution of the mammalian middle ear and jaw: Adaptations and novel structures. *Journal of Anatomy*, 222(1), 147–160. <https://doi.org/10.1111/j.1469-7580.2012.01526.x>

Bakkers, J., Hild, M., Kramer, C., Furutani-Seiki, M., & Hammerschmidt, M. (2002). Zebrafish Δ Np63 is a direct target of bmp signaling and encodes a transcriptional repressor blocking neural specification in the ventral ectoderm. *Developmental Cell*, 2(5), 617–627. [https://doi.org/10.1016/S1534-5807\(02\)00163-6](https://doi.org/10.1016/S1534-5807(02)00163-6)

Ben, J., Jabs, E. W., & Chong, S. S. (2005). Genomic, cDNA and embryonic expression analysis of zebrafish IRF6, the gene mutated in the human oral clefting disorders Van der Woude and popliteal pterygium syndromes. *Gene Expression Patterns*, 5(5), 629–638. <https://doi.org/10.1016/j.modgep.2005.03.002>

Boulet, S. L., Rasmussen, S. A., & Honein, M. A. (2008). A population-based study of Craniosynostosis in metropolitan Atlanta, 1989–2003. *American Journal of Human Genetics*, 146A(8), 984–991. <https://doi.org/10.1002/ajmg.a.32208>

Braybrook, C., Lisgo, S., Doudney, K., Henderson, D., Marciano, A., Strachan, T., ... Lindsay, S. (2002). Craniofacial expression of human and murine TBX22 correlates with the cleft palate and ankyloglossia phenotype observed in CPX patients. *Human Molecular Genetics*, 11(22), 2793–2804. <https://doi.org/10.1093/hmg/11.22.2793>

Burdick, A., Bixler, D., & Puckett, C. (1985). Genetic analysis in families with Van Der Woude syndrome. *Journal of Craniofacial Genetics and Developmental Biology*, 5(2), 181–208.

Bush, J. O., & Jiang, R. (2012). Palatogenesis: Morphogenetic and molecular mechanisms of secondary palate development. *Development*, 139(2), 231–243. <https://doi.org/10.1242/dev.067082>

Byvaltsev, V., Belykh, O., & Belykh, E. (2012). New aspects in the epidemiology of craniofacial anomalies. *World Neurosurgery*, 77(5–6), 599–600. <https://doi.org/10.1016/j.wneu.2012.03.004>

Celli, J., Duijff, P., Hamel, B. C. J., Bamshad, M., Kramer, B., Smits, A. P. T., ... van Bokhoven, H. (1999). Heterozygous germline mutations in the p53 homolog p63 are the cause of EEC syndrome. *Cell*, 99, 143–153. [https://doi.org/10.1016/S0092-8674\(00\)81646-3](https://doi.org/10.1016/S0092-8674(00)81646-3)

Chotzen, F. (1932). Eine eigenartige familiäre Entwicklungsstörung (Akrocephalosyndaktylie, Dysostosis craniofacialis und Hyper-telorismus). *Msschr Kinderheilk*, 55, 97–122.

Clouthier, D. E., Garcia, E., & Schilling, T. F. (2010). Regulation of facial morphogenesis by endothelin signaling: Insights from mice and fish. *American Journal of Medical Genetics. Part A*, 152A(12), 2962–2973. <https://doi.org/10.1002/ajmg.a.33568>

Clouthier, D. E., Hosoda, K., Richardson, J. A., Williams, S. C., Yanagisawa, H., Kuwaki, Y., ... Yanagisawa, M. (1998). Cranial and cardiac neural crest defects in endothelin-a receptor-deficient mice. *Development*, 125, 813–824. <https://doi.org/10.1242/dev.00339>

Connerney, J., Andreeva, V., Leshem, Y., Mercado, M. A., Dowell, K., Yang, X., ... Spicer, D. B. (2008). Twist1 homodimers enhance FGF responsiveness of the cranial sutures and promote suture closure. *Developmental Biology*, 318(2), 323–334. <https://doi.org/10.1016/j.ydbio.2008.03.037>

Das, A., & Crump, J. G. (2012). Bmps and id2a act upstream of Twist1 to restrict ectomesenchyme potential of the cranial neural crest. *PLoS Genetics*, 8(5), e1002710. <https://doi.org/10.1371/journal.pgen.1002710>

Davis, C. A., Holmyard, D. P., Millen, K. J., & Joyner, A. L. (1991). Examining pattern formation in mouse, chicken and frog embryos with an en-specific antiserum. *Development*, 111, 287–298. Retrieved from <https://dev.biologists.org/content/111/2/287.long>

de la Garza, G., Schleiffarth, J. R., Dunnwald, M., Mankad, A., Weirather, J. L., Bonde, G., ... Cornell, R. A. (2013). Interferon regulatory factor 6 promotes differentiation of the periderm by activating expression of Grainyhead-like 3. *The Journal of Investigative Dermatology*, 133(1), 68–77. <https://doi.org/10.1038/jid.2012.269>

Dougherty, M., Kamel, G., Grimaldi, M., Grerer, L., Shubinets, V., Ethier, R., ... Liao, E. C. (2013). Distinct requirements for wnt9a and irf6 in extension and integration mechanisms during zebrafish palate morphogenesis. *Development*, 140(1), 76–81. <https://doi.org/10.1242/dev.080473>

Duncan, K. M., Mukherjee, K., Cornell, R. A., & Liao, E. C. (2017). Zebrafish models of orofacial clefts. *Developmental Dynamics*, 246(11), 897–914. <https://doi.org/10.1002/dvdy.24566>

Dworkin, S., Simkin, J., Darido, C., Partridge, D. D., Georgy, S. R., Caddy, J., ... Jane, S. M. (2014). Grainyhead-like 3 regulation of endothelin-1 in the pharyngeal endoderm is critical for growth and development of the craniofacial skeleton. *Mechanisms of Development*, 133, 77–90. <https://doi.org/10.1016/j.mod.2014.05.005>

Eberhart, J. K., He, X., Swartz, M. E., Yan, Y.-L., Song, H., Boling, T. C., ... Postlethwait, J. (2008). MicroRNA mir140 modulates Pdgf signaling during palatogenesis. *Nature Genetics*, 40(3), 290–298. <https://doi.org/10.1038/ng.82>

el Ghouzzi, V., Le Merrer, M., Perrin-Schmitt, F., Lajeunie, E., Benit, P., Renier, D., ... Bonaventure, J. (1997). Mutations of the TWIST gene in

- the Saethre-Chotzen syndrome. *Nature Genetics*, 15(1), 42–46. <https://doi.org/10.1038/ng0197-42>
- Erickson, C. (1985). Control of neural crest cell dispersion in the trunk of the avian embryo. *Developmental Biology*, 111, 138–157. [https://doi.org/10.1016/0012-1606\(85\)90442-7](https://doi.org/10.1016/0012-1606(85)90442-7)
- Fakhouri, W. D., Metwalli, K., Naji, A., Bakhiet, S., Quispe-Salcedo, A., Nitschke, L., ... Schutte, B. C. (2017). Intracellular genetic interaction between *Irf6* and *Twist1* during craniofacial Development. *Scientific Reports*, 7, 1–14. <https://doi.org/10.1038/s41598-017-06310-z>
- Ferrante, M. I., Romio, L., Castro, S., Collins, J. E., Goulding, D. A., Stemple, D. L., ... Wilson, S. W. (2009). Convergent extension movements and ciliary function are mediated by *ofd1*, a zebrafish orthologue of the human oral-facial-digital type 1 syndrome gene. *Human Molecular Genetics*, 18(2), 289–303. <https://doi.org/10.1093/hmg/ddn356>
- Fuller, K., O'Connell, J. T., Gordon, J., Mauti, O., & Eggenschwiler, J. (2014). Rab23 regulates nodal signaling in vertebrate left-right patterning independently of the hedgehog pathway. *Developmental Biology*, 391(2), 182–195. <https://doi.org/10.1016/j.ydbio.2014.04.012>
- Gardner, C. A., & Barald, K. F. (1992). Expression of engrailed-like proteins in the chick embryo. *Developmental Dynamics*, 193(4), 370–388. <https://doi.org/10.1002/aja.1001930410>
- Gegenbaur, C. (1878). *Elements of comparative anatomy*. London, UK: Macmillan.
- Gillis, J. A., Dahn, R. D., & Shubin, N. H. (2009). Shared developmental mechanisms pattern the vertebrate gill arch and paired fin skeletons. *Proceedings of the National Academy of Sciences of the United States of America*, 106(14), 5720–5724. <https://doi.org/10.1073/pnas.0810959106>
- Gillis, J. A., & Hall, B. K. (2016). A shared role for sonic hedgehog signalling in patterning chondrichthyan gill arch appendages and tetrapod limbs. *Development*, 143(8), 1313–1317. <https://doi.org/10.1242/dev.133884>
- Global Registry and Database on Craniofacial Anomalies. (2001). Retrieved from Barau, Brazil. <https://www.who.int/genomics/anomalies/en/CFA-RegistryMeeting-2001.pdf>
- Grandel, H., Lun, K., Rauch, G.-J., Rhinn, M., Piotrowski, T., Houart, C., ... Brand, M. (2002). Retinoic acid signalling in the zebrafish embryo is necessary during pre-segmentation stages to pattern the anterior-posterior axis of the CNS and to induce a pectoral fin bud. *Development*, 129(12), 2851–2865. Retrieved from <https://dev.biologists.org/content/develop/129/12/2851.full.pdf>
- Grandel, H., & Schulte-Merker, S. (1998). The development of the paired fins in the Zebrafish (*Danio rerio*). *Mechanisms of Development*, 79, 99–120. [https://doi.org/10.1016/S0925-4773\(98\)00176-2](https://doi.org/10.1016/S0925-4773(98)00176-2)
- Grim, M., & Christ, B. (1993). Neural crest cell migration into the limb bud of avian embryos. *Progress in Clinical and Biological Research*, 383A, 391–402. Retrieved from <https://pubmed.ncbi.nlm.nih.gov/7508130/>
- Hilliard, S., Yu, L., Gu, S., Zhang, Z., & Chen, Y. (2005). Regional regulation of palatal growth and patterning along the anterior-posterior axis in mice. *Journal of Anatomy*, 207(5), 655–667. <https://doi.org/10.1111/j.1469-7580.2005.00474.x>
- Hirsch, N., Eshel, R., Bar Yaacov, R., Shahar, T., Shmulevich, F., Dahan, I., ... Birnbaum, R. Y. (2018). Unraveling the transcriptional regulation of TWIST1 in limb development. *PLoS Genetics*, 14(10), e1007738. <https://doi.org/10.1371/journal.pgen.1007738>
- Holmes, G. (2012). The role of vertebrate models in understanding Craniosynostosis. *Child's Nervous System*, 28(9), 1471–1481. <https://doi.org/10.1007/s00381-012-1844-3>
- Howard, T., Paznekas, W., Green, E., Chiang, L., Ma, N., Ortiz de Luna, R., ... Jabs, E. W. (1997). Mutations in TWIST, a basic helix-loop-helix transcription factor, in Saethre-Chotzen syndrome. *Nature Genetics*, 15, 36–41. <https://doi.org/10.1038/ng0197-36>
- Ingraham, C. R., Kinoshita, A., Kondo, S., Yang, B., Sajan, S., Trout, K. J., ... Schutte, B. C. (2006). Abnormal skin, limb and craniofacial morphogenesis in mice deficient for interferon regulatory factor 6 (*Irf6*). *Nature Genetics*, 38(11), 1335–1340. <https://doi.org/10.1038/ng1903>
- Kague, E., Gallagher, M., Burke, S., Parsons, M., Franz-Odenaal, T., & Fisher, S. (2012). Skeletogenic fate of Zebrafish cranial and trunk neural crest. *PLoS One*, 7(11), 1–13. <https://doi.org/10.1371/journal.pone.0047394>
- Kantaputra, P. N., & Carlson, B. M. (2019). Genetic regulatory pathways of split-hand/foot malformation. *Clinical Genetics*, 95(1), 132–139. <https://doi.org/10.1111/cge.13434>
- Kashgari, G., Meinecke, L., Gordon, W., Ruiz, B., Yang, J., Ma, A. L., ... Andersen, B. (2020). Epithelial migration and non-adhesive periderm are required for digit separation during mammalian Development. *Developmental Cell*, 52(6), 764–778. <https://doi.org/10.1016/j.devcel.2020.01.032>
- Kawata, M., Taniguchi, Y., Mori, D., Yano, F., Ohba, S., Chung, U.-I., ... Saito, T. (2017). Different regulation of limb development by p63 transcript variants. *PLoS One*, 12(3), 1–19. <https://doi.org/10.1371/journal.pone.0174122>
- Kawauchi, S., Santos, R., Muto, A., Lopez-Burks, M. E., Schilling, T. F., Lander, A. D., & Calof, A. L. (2016). Using mouse and zebrafish models to understand the etiology of developmental defects in Cornelia de Lange syndrome. *American Journal of Medical Genetics. Part C, Seminars in Medical Genetics*, 172(2), 138–145. <https://doi.org/10.1002/ajmg.c.31484>
- Kimmel, C. B., Ullmann, B., Walker, M., Miller, C. T., & Crump, J. G. (2003). Endothelin 1-mediated regulation of pharyngeal bone development in zebrafish. *Development*, 130(7), 1339–1351. <https://doi.org/10.1242/dev.00338>
- Kondo, S., Schutte, B. C., Richardson, R. J., Bjork, B. C., Knight, A. S., Watanabe, Y., ... Murray, J. C. (2002). Mutations in IRF6 cause Van der Woude and popliteal pterygium syndromes. *Nature Genetics*, 32(2), 285–289. <https://doi.org/10.1038/ng985>
- Kurihara, Y., H, K., Suzuki, H., Kodama, T., Maemura, K., Nagai, R., ... Yazaki, Y. (1994). Elevated blood pressure and craniofacial abnormalities in mice deficient in endothelin-1. *Nature*, 368, 703–710.
- Larbuissou, A., Dalcq, J., Martial, J. A., & Muller, M. (2013). Fgf receptors Fgfr1a and Fgfr2 control the function of pharyngeal endoderm in late cranial cartilage development. *Differentiation*, 86(4–5), 192–206. <https://doi.org/10.1016/j.diff.2013.07.006>
- Laufer, E., Nelson, C. E., Johnson, R. L., Morgan, B. A., & Tabin, C. (1994). Sonic hedgehog and Fgf-4 act through a signaling cascade and feedback loop to integrate growth and patterning of the developing limb bud. *Cell*, 79(6), 993–1003. [https://doi.org/10.1016/0092-8674\(94\)90030-2](https://doi.org/10.1016/0092-8674(94)90030-2)
- Lee, H., & Kimelman, D. (2002). A dominant-negative form of p63 is required for epidermal proliferation in zebrafish. *Developmental Cell*, 2, 607–616. [https://doi.org/10.1016/S1534-5807\(02\)00166-1](https://doi.org/10.1016/S1534-5807(02)00166-1)
- Leerberg, D. M., Hopton, R. E., & Draper, B. W. (2019). Fibroblast growth factor receptors function redundantly during zebrafish embryonic development. *Genetics*, 212(4), 1301–1319. <https://doi.org/10.1534/genetics.119.302345>
- Leong, I. U., Skinner, J. R., Shelling, A. N., & Love, D. R. (2013). Expression of a mutant *kcnj2* gene transcript in zebrafish. *ISRN Molecular Biology*, 2013, 324839. <https://doi.org/10.1155/2013/324839>
- Letelier, J., de la Calle-Mustienes, E., Pieretti, J., Naranjo, S., Maeso, I., Nakamura, T., ... Gomez-Skarmeta, J. L. (2018). A conserved Shh cis-regulatory module highlights a common developmental origin of unpaired and paired fins. *Nature Genetics*, 50(4), 504–509. <https://doi.org/10.1038/s41588-018-0080-5>
- Lo Iacono, N., Mantero, S., Chiarelli, A., Garcia, E., Mills, A. A., Morasso, M. I., ... Merlo, G. R. (2008). Regulation of Dlx5 and Dlx6 gene expression by p63 is involved in EEC and SHFM congenital limb defects. *Development*, 135(7), 1377–1388. <https://doi.org/10.1242/dev.011759>

- Loomis, C. A., Kimmel, R. A., Tong, C.-X., Michaud, J., & Joyner, A. L. (1998). Analysis of the genetic pathway leading to formation of ectopic apical ectodermal ridges in mouse *Engrailed-1* mutant limbs. *Development*, 125, 1137–1148. Retrieved from. <https://dev.biologists.org/content/125/6/1137.article-info>
- Lorent, K., Yeo, S. Y., Oda, T., Chandrasekharappa, S., Chitnis, A., Matthews, R. P., & Pack, M. (2004). Inhibition of jagged-mediated notch signaling disrupts zebrafish biliary development and generates multi-organ defects compatible with an Alagille syndrome phenocopy. *Development*, 131(22), 5753–5766. <https://doi.org/10.1242/dev.01411>
- Mercader, N. (2007). Early steps of paired fin development in zebrafish compared with tetrapod limb development. *Development, Growth & Differentiation*, 49(6), 421–437. <https://doi.org/10.1111/j.1440-169X.2007.00942.x>
- Merrill, A. E., Bochukova, E. G., Brugger, S. M., Ishii, M., Pilz, D. T., Wall, S. A., ... Maxson, R. E., Jr. (2006). Cell mixing at a neural crest-mesoderm boundary and deficient ephrin-Eph signaling in the pathogenesis of craniosynostosis. *Human Molecular Genetics*, 15(8), 1319–1328. <https://doi.org/10.1093/hmg/ddl052>
- Miles, L. B., Darido, C., Kaslin, J., Heath, J. K., Jane, S. M., & Dworkin, S. (2017). Mis-expression of grainyhead-like transcription factors in zebrafish leads to defects in enveloping layer (EVL) integrity, cellular morphogenesis and axial extension. *Scientific Reports*, 7(1), 17607. <https://doi.org/10.1038/s41598-017-17898-7>
- Miller, C. T., Schilling, T. F., Lee, K., Parker, J., & Kimmel, C. B. (2000). Sucker encodes a zebrafish endothelin-1 required for ventral pharyngeal arch development. *Development*, 127(17), 3815–3828. Retrieved from. <https://dev.biologists.org/content/development/127/17/3815.full.pdf>
- Miller, C. T., Swartz, M. E., Khuu, P. A., Walker, M. B., Eberhart, J. K., & Kimmel, C. B. (2007). *mef2ca* is required in cranial neural crest to effect Endothelin1 signaling in zebrafish. *Developmental Biology*, 308(1), 144–157. <https://doi.org/10.1016/j.ydbio.2007.05.018>
- Miller, C. T., Yelon, D., Stainier, D. Y. R., & Kimmel, C. B. (2003). Two *endothelin 1* effectors, *hand2* and *bapx1*, pattern ventral pharyngeal cartilage and the jaw joint. *Development*, 130, 1353–1365. <https://doi.org/10.1242/dev.00339>
- Miyashita, T., Baddam, P., Smeeton, J., Oel, A. P., Natarajan, N., Gordon, B., Palmer, A. R., Crump, J. G., Graf, D. Allison, W. T. (2020). *nkx3.2* mutant zebrafish accommodate jaw joint loss through a phenocopy of the head shapes of Paleozoic jawless fish. *The Journal of Experimental Biology*, 223(Pt 15). <https://doi.org/10.1242/jeb.216945>
- Monnich, M., Kuriger, Z., Print, C. G., & Horsfield, J. A. (2011). A zebrafish model of Roberts syndrome reveals that *Esco2* depletion interferes with development by disrupting the cell cycle. *PLoS One*, 6(5), e20051. <https://doi.org/10.1371/journal.pone.0020051>
- Mork, L., & Crump, G. (2015). Zebrafish craniofacial Development: A window into early patterning. *Current Topics in Developmental Biology*, 115, 235–269. <https://doi.org/10.1016/bs.ctdb.2015.07.001>
- Morris-Kay, G. M. (2001). Derivation of the mammalian skull vault. *Journal of Anatomy*, 199, 143–151. <https://doi.org/10.1046/j.1469-7580.2001.19910143.x>
- Ng, J. K., Kawakami, Y., Buscher, D., Raya, A., Itoh, T., Koth, C. M., ... Izpisua Belmonte, J. C. (2002). The limb identity gene *Tbx5* promotes limb initiation by interacting with *Wnt2b* and *Fgf10*. *Development*, 129, 5161–5170. Retrieved from. <http://dev.biologists.org/content/129/22/5161>
- Nie, X., Luukko, K., & Kettunen, P. (2006). FGF signaling in craniofacial development and developmental disorders. *Oral Diseases*, 12(2), 102–111. <https://doi.org/10.1111/j.1601-0825.2005.01176.x>
- Niswander, L., Jeffrey, S., Martin, G. R., & Tickle, C. (1994). A positive feedback loop coordinates growth and patterning in the vertebrate limb. *Nature*, 371, 609–612. <https://doi.org/10.1038/371609a0>
- Niswander, L., & Martin, G. R. (1993). FGF-4 and BMP-2 have opposite effects on limb growth. *Nature*, 361(6407), 68–71. <https://doi.org/10.1038/361068a0>
- Niswander, L., Tickle, C., Vogel, A., Booth, I., & Martin, G. R. (1993). FGF-4 replaces the apical ectodermal ridge and directs outgrowth and patterning of the limb. *Cell*, 75(3), 579–587. [https://doi.org/10.1016/0092-8674\(93\)90391-3](https://doi.org/10.1016/0092-8674(93)90391-3)
- Ohuchi, H., Nakagawa, T., Yamamoto, A., Araga, A., Ohata, T., Ishimaru, T., ... Noji, S. (1997). The mesenchymal factor, FGF10, initiates and maintains the outgrowth of the chick limb bud through interaction with FGF8, an apical ectodermal factor. *Development*, 124, 2235–2244. Retrieved from. <http://dev.biologists.org/content/124/11/2235.article-info>
- O'Rourke, M., Soo, K., Behringer, R. R., Hui, C.-C., & Tam, P. (2002). Twist plays an essential role in FGF and SHH signal transduction during mouse development. *Developmental Biology*, 248(1), 143–156. <https://doi.org/10.1006/dbio.2002.0730>
- Pajni-Underwood, S., Wilson, C. P., Elder, C., Mishina, Y., & Lewandoski, M. (2007). BMP signals control limb bud interdigital programmed cell death by regulating FGF signaling. *Development*, 134(12), 2359–2368. <https://doi.org/10.1242/dev.001677>
- Panthaki, Z. J., & Armstrong, M. B. (2003). Hand abnormalities associated with craniofacial syndromes. *Journal of Craniofacial Surgery*, 14(5), 709–712. <https://doi.org/10.1097/00001665-200309000-00020>
- Parr, B. A., & McMahon, A. P. (1995). Dorsalizing signal *Wnt-7a* required for normal polarity of D-V and a-P axes of mouse limb. *Nature*, 374, 350–353. <https://doi.org/10.1038/374350a0>
- Paznekas, W., Cunningham, M. L., Howard, T., Korf, B., Lipson, M., Grix, A., ... Jabs, E. W. (1998). Genetic heterogeneity of Saethre-Chotzen syndrome, due to *TWIST* and *FGFR* mutations. *AJHG*, 62(6), 1370–1380. <https://doi.org/10.1086/301855>
- Penchaszadeh, V. B., & de Negrotti, T. C. (1976). Ectrodactyly-ectodermal dysplasia-clefting (EEC) syndrome: Dominant inheritance and variable expression. *Journal of Medical Genetics*, 13(4), 381–384. <https://doi.org/10.1136/jmg.13.4.281>
- Percival, S. M., Thomas, H. R., Amsterdam, A., Carroll, A. J., Lees, J. A., Yost, H. J., & Parant, J. M. (2015). Variations in dysfunction of sister chromatid cohesion in *esco2* mutant zebrafish reflect the phenotypic diversity of Roberts syndrome. *Disease Models & Mechanisms*, 8(8), 941–955. <https://doi.org/10.1242/dmm.019059>
- Person, A. D., Beiraghi, S., Sieben, C. M., Hermanson, S., Neumann, A. N., Robu, M. E., ... Lohr, J. L. (2010). *WNT5A* mutations in patients with autosomal dominant Robinow syndrome. *Developmental Dynamics*, 239(1), 327–337. <https://doi.org/10.1002/dvdy.22156>
- Peters, H., Neubuser, A., Kratochwil, K., & Balling, R. (1998). *Pax9*-deficient mice lack pharyngeal pouch derivatives and teeth and exhibit craniofacial and limb abnormalities. *Genes & Development*, 12(17), 2735–2747. <https://doi.org/10.1101/gad.12.17.2735>
- Petit, F., Sears, K. E., & Ahituv, N. (2017). Limb development: A paradigm of gene regulation. *Nature*, 18, 245–258. <https://doi.org/10.1038/nrg.2016.167>
- Peyrard-Janvid, M., Leslie, E., Kousa, Y., Smith, T., Dunnwald, M., Magnusson, M., ... Schutte, B. (2014). Dominant mutations in *GRHL3* cause Van Der Woude syndrome and disrupt Oral periderm Development. *American Journal of Human Genetics*, 94(1), 23–32. <https://doi.org/10.1016/j.ajhg.2013.11.009>
- Pistocchi, A., Fazio, G., Cereda, A., Ferrari, L., Bettini, L. R., Messina, G., ... Massa, V. (2013). Cornelia de Lange syndrome: NIPBL haploinsufficiency downregulates canonical Wnt pathway in zebrafish embryos and patients fibroblasts. *Cell Death & Disease*, 4, e866. <https://doi.org/10.1038/cddis.2013.371>
- Pizette, S., & Niswander, L. (1999). BMPs negatively regulate structure and function of the limb apical ectodermal ridge. *Development*, 126, 883–894. Retrieved from. <https://dev.biologists.org/content/126/5/883.long>

- Quarto, N., & Longaker, M. T. (2005). The zebrafish (*Danio rerio*): A model system for cranial suture patterning. *Cells, Tissues, Organs*, 181, 109–118. <https://doi.org/10.1159/000091100>
- Riddle, R. D., Johnson, R. L., Laufer, E., & Tabin, C. (1993). *Sonic hedgehog* mediates the polarizing activity of the ZPA. *Cell*, 75, 1401–1416. [https://doi.org/10.1016/0092-8674\(93\)90626-2](https://doi.org/10.1016/0092-8674(93)90626-2)
- Roberts, J. B. (1919). A child with double cleft of lip and palate, protrusion of the intermaxillary portion of the upper jaw and imperfect development of the bones of the four extremities. *Annals of surgery*, 70, 252–234.
- Rolef Ben-Shahar, T., Heeger, S., Lehane, C., East, P., Flynn, H., Skehel, M., & Uhlmann, F. (2008). Eco-1 dependent cohesin acetylation during establishment of sister chromatid cohesion. *Science*, 321 (5888), 563–566. <https://doi.org/10.1126/science.1157774>
- Sabel, J. L., d'Alencon, C., O'Brien, E. K., Van Otterloo, E., Lutz, K., Cuykendall, T. N., ... Cornell, R. A. (2009). Maternal interferon regulatory factor 6 is required for the differentiation of primary superficial epithelia in *Danio* and *Xenopus* embryos. *Developmental Biology*, 325 (1), 249–262. <https://doi.org/10.1016/j.ydbio.2008.10.031>
- Saethre, H. (1931). Ein Beitrag zum Turmschädelproblem, (Pathogenese, Erblichkeit und Symptomatologie). *Deutsche Zeitschrift f. Nervenheilkunde*, 117, 533–555. <https://doi.org/10.1007/BF01673869>
- Santagati, F., & Rijli, F. M. (2003). Cranial neural crest and the building of the vertebrate head. *Nature Reviews. Neuroscience*, 4(10), 806–818. <https://doi.org/10.1038/nrn1221>
- Santos-Pereira, J. M., Gallardo-Fuentes, L., Neto, A., Acemel, R. D., & Tena, J. J. (2019). Pioneer and repressive functions of p63 during zebrafish embryonic ectoderm specification. *Nature Communications*, 10(1), 3049. <https://doi.org/10.1038/s41467-019-11121-z>
- Saunders, J. W. (1948). The proximo-distal sequence of the origin of the parts of the chick wing and the role of the ectoderm. *The Journal of Experimental Zoology*, 108, 363–404. <https://doi.org/10.1002/jez.1401080304>
- Saunders, J. W., & Gasseling, M. T. (1968). In R. Fleischmajer & R. F. Billingham (Eds.), *Ectodermal-mesenchymal interactions in the origin of limb symmetry*. Baltimore: Williams & Wilkins.
- Schilling, T. F., & Kimmel, C. B. (1997). Musculoskeletal patterning in the pharyngeal segments of the zebrafish embryo. *Development*, 124, 2945–2960. Retrieved from. <https://dev.biologists.org/content/124/15/2945>
- Sekine, K., Ohuchi, H., Fujiwara, M., Yamasaki, M., Yoshizawa, T., Sato, T., ... Kato, S. (1999). Fgf10 is essential for limb and lung formation. *Nature Genetics*, 21, 138–141. <https://doi.org/10.1038/5096>
- Selever, J., Liu, W., Lu, M., Behringer, R. R., & Martin, J. F. (2004). Bmp4 in limb bud mesoderm regulates digit pattern by controlling AER development. *Developmental Biology*, 276(2), 268–279. <https://doi.org/10.1016/j.ydbio.2004.08.024>
- Serrano, M., Demarest, B. L., Tone-Pah-Hote, T., Tristani-Firouzi, M., & Yost, J. (2019). Inhibition of notch signaling rescues cardiovascular development in kabuki syndrome. *PLoS Biology*, 17(9), e3000087. <https://doi.org/10.1371/journal.pbio.3000087>
- Sleight, V. A., & Gillis, J. A. (2020). Embryonic origin and serial homology of gill arches and paired fins in the skate, *Leucoraja erinacea*. *eLife*, 9 (e60635), 1–14. <https://doi.org/10.7554/eLife.60635>
- Smith, J. D., Hing, A. V., Clarke, C. M., Johnson, N. M., Perez, F. A., Park, S. S., ... Cunningham, M. L. (2014). Exome sequencing identifies a recurrent de novo ZSWIM6 mutation associated with acromelic frontonasal dysostosis. *American Journal of Human Genetics*, 95(2), 235–240. <https://doi.org/10.1016/j.ajhg.2014.07.008>
- Sun, X., Mariani, F. V., & Martin, G. R. (2002). Functions of FGF signalling from the apical ectodermal ridge in limb development. *Nature*, 418, 501–508. <https://doi.org/10.1038/nature00902>
- Swartz, M. E., Sheehan-Rooney, K., Dixon, M. J., & Eberhart, J. K. (2011). Examination of a palatogenic gene program in zebrafish. *Developmental Dynamics*, 240(9), 2204–2220. <https://doi.org/10.1002/dvdy.22713>
- Szabo-Rogers, H. L., Smithers, L. E., Yakob, W., & Liu, K. J. (2010). New directions in craniofacial morphogenesis. *Developmental Biology*, 341 (1), 84–94. <https://doi.org/10.1016/j.ydbio.2009.11.021>
- Teng, C. S., Ting, M. C., Farmer, D. T., Brockop, M., Maxson, R. E., & Crump, J. G. (2018). Altered bone growth dynamics prefigure cranio-synostosis in a zebrafish model of Saethre-Chotzen syndrome. *eLife*, 7(e37024), 1–23. <https://doi.org/10.7554/eLife.37024>
- Topczewska, J. M., Shoela, R. A., Tomaszewski, J. P., Mirmira, R. B., & Gosain, A. K. (2016). The morphogenesis of cranial sutures in zebrafish. *PLoS One*, 11(11), e0165775. <https://doi.org/10.1371/journal.pone.0165775>
- Twigg, S. R., Lloyd, D., Jenkins, D., Elcioglu, N. E., Cooper, C. D., Al-Sannaa, N., ... Wilkie, A. O. (2012). Mutations in multidomain protein MEGF8 identify a carpenter syndrome subtype associated with defective lateralization. *American Journal of Human Genetics*, 91(5), 897–905. <https://doi.org/10.1016/j.ajhg.2012.08.027>
- Twigg, S. R., & Wilkie, A. O. (2015). New insights into craniofacial malformations. *Human Molecular Genetics*, 24(R1), R50–R59. <https://doi.org/10.1093/hmg/ddv228>
- Tyurina, O. V., Guner, B., Popova, E., Feng, J., Schier, A. F., Kohtz, J. D., & Karlstrom, R. O. (2005). Zebrafish Gli3 functions as both an activator and a repressor in hedgehog signaling. *Developmental Biology*, 277(2), 537–556. <https://doi.org/10.1016/j.ydbio.2004.10.003>
- Update on Overall Prevalence of Major Birth Defects. (2008). Retrieved from Atlanta, Georgia. <https://www.cdc.gov/mmwr/preview/mmwrhtml/mm5701a2.htm>
- van Eeden, F. J. M., Granato, M., Schach, U., Brand, M., Furutani-Seiki, M., Haffter, P., ... Nüsslein-Volhard, C. (1996). Genetic analysis of fin formation in the zebrafish, *Danio rerio*. *Development*, 123, 255–262. Retrieved from <https://dev.biologists.org/content/develop/123/1/255.full.pdf>
- Van Laarhoven, P. M., Neitzel, L. R., Quintana, A. M., Geiger, E. A., Zackai, E. H., Clouthier, D. E., ... Shaikh, T. H. (2015). Kabuki syndrome genes KMT2D and KDM6A: Functional analyses demonstrate critical roles in craniofacial, heart and brain development. *Human Molecular Genetics*, 24(15), 4443–4453. <https://doi.org/10.1093/hmg/ddv180>
- Vasluiu, E., van der Sluis, C. K., van Essen, A. J., Bergman, J. E. H., Dijkstra, P. U., Reinders-Messelink, H. A., & de Walle, H. E. K. (2013). Birth prevalence for congenital limb defects in the northern Netherlands: A 30-year population-based study. *BMC Musculoskeletal Disorders*, 14(323), 1–14. <https://doi.org/10.1186/1471-2474-14-323>
- Vega, H., Waisfisz, Q., Gordillo, M., Sakai, N., Yanagihara, I., Yamada, M., ... Joenje, H. (2005). Roberts syndrome is caused by mutations in ESCO2, a human homolog of yeast ECO1 that is essential for the establishment of sister chromatid cohesion. *Nature Genetics*, 37, 468–470. <https://doi.org/10.1038/ng1548>
- Wada, N., Javidan, Y., Nelson, S., Carney, T. J., Kelsh, R. N., & Schilling, T. F. (2005). Hedgehog signaling is required for cranial neural crest morphogenesis and chondrogenesis at the midline in the zebrafish skull. *Development*, 132, 3977–3988. <https://doi.org/10.1242/dev.01943>
- Walshe, J., & Mason, I. (2003). Fgf signaling is required for formation of cartilage in the head. *Developmental Biology*, 264, 522–536. <https://doi.org/10.1016/j.ydbio.2003.08.010>
- Weaver, K. N., Watt, K. E., Hufnagel, R. B., Navajas Acedo, J., Linscott, L. L., Sund, K. L., ... Saal, H. M. (2015). Acrofacial dysostosis, cincinnati type, a mandibulofacial dysostosis syndrome with limb anomalies, is caused by POLR1A dysfunction. *American Journal of Human Genetics*, 96(5), 765–774. <https://doi.org/10.1016/j.ajhg.2015.03.011>
- Wenger, T., Miller, D., & Evans, K. (1998–2020). FGFR Craniosynostosis syndromes overview. In M. Adam, H. Ardinger, R. Pagon, & S. Wallace

- (Eds.), *GeneReviews*® [Internet]. Seattle, WA: GeneReviews. Retrieved from <https://www.ncbi.nlm.nih.gov/books/NBK1455/>
- Wilcox, W. R., Coulter, C. P., & Schmitz, M. L. (2015). Congenital limb deficiency disorders. *Clinics in Perinatology*, 42(2), 281–300, viii. <https://doi.org/10.1016/j.clp.2015.02.004>
- Wilson, J., & Tucker, A. S. (2004). Fgf and bmp signals repress the expression of *Bapx1* in the mandibular mesenchyme and control the position of the developing jaw joint. *Developmental Biology*, 266, 138–150. <https://doi.org/10.1016/j.ydbio.2003.10.012>
- Wu, X., & Gu, Y. (2019). Signaling mechanisms underlying genetic pathophysiology of craniosynostosis. *International Journal of Biological Sciences*, 15(2), 298–311. <https://doi.org/10.7150/ijbs.29183>
- Yang, A., Kaghad, M., Wang, Y., Gillett, E., Fleming, M., Dötsch, V., ... McKeon, F. (1998). *p63*, a *p53* homolog at 3q27-29, encodes multiple products with transactivating, death-inducing, and dominant-negative activities. *Molecular Cell*, 2(3), 305–316. [https://doi.org/10.1016/S1097-2765\(00\)80275-0](https://doi.org/10.1016/S1097-2765(00)80275-0)
- Yang, A., Schweitzer, R., Sun, D., Kaghad, M., Walker, N., Bronson, R., ... McKeon, F. (1999). *p63* is essential for regenerative proliferation in limb, craniofacial, and epithelial development. *Nature*, 398, 714–718. <https://doi.org/10.1038/19539>
- Yano, T., & Tamura, K. (2013). The making of differences between fins and limbs. *Journal of Anatomy*, 222(1), 100–113. <https://doi.org/10.1111/j.1469-7580.2012.01491.x>
- Zuniga, A. (2015). Next generation limb development and evolution: Old questions, new perspectives. *Development*, 142(22), 3810–3820. <https://doi.org/10.1242/dev.125757>
- Zuniga, E., Stellabotte, F., & Crump, J. G. (2010). Jagged-notch signaling ensures dorsal skeletal identity in the vertebrate face. *Development*, 137(11), 1843–1852. <https://doi.org/10.1242/dev.049056>

How to cite this article: Truong BT, Artinger KB. The power of zebrafish models for understanding the co-occurrence of craniofacial and limb disorders. *genesis*. 2021;1–19. <https://doi.org/10.1002/dvg.23407>